

New characterizations, in the framework of critical kernels, of 2D, 3D and 4D minimal non-simple sets and P-simple points

Gilles Bertrand and Michel Couprie
Université Paris-Est, LABINFO-IGM,
UMR CNRS 8049, A2SI-ESIEE, France
e-mail: (m.couprie,g.bertrand)@esiee.fr

Abstract

Critical kernels constitute a general framework in the category of abstract complexes for the study of parallel homotopic thinning in any dimension. In this article, we present new results linking critical kernels to minimal non-simple sets (MNS) and P-simple points, which are notions conceived to study parallel thinning in discrete grids. We show that these two previously introduced notions can be retrieved, better understood and enriched in the framework of critical kernels. In particular, we propose new characterizations which hold in dimensions 2, 3 and 4, and which lead to efficient algorithms for detecting P-simple points and minimal non-simple sets.

Key Words: Parallel thinning, topology preservation, critical kernel, P-simple point, minimal non-simple set, cubical complex, collapse, simple point, confluence, 4D space.

Introduction

Topology-preserving operators, such as homotopic skeletonization, are used in many applications of image analysis to transform an object while leaving unchanged its topological characteristics. In discrete grids (\mathbb{Z}^2 , \mathbb{Z}^3 , \mathbb{Z}^4), such a transformation can be defined thanks to the notion of simple point [21, 13]: intuitively, a point (or pixel in 2D, voxel in 3D) of an object is called simple if it can be deleted from this object without altering its topology. See for example Fig. 1a, where simple points are shown in gray.

The most “natural” way to thin an object consists of removing some of its border points in parallel, in a symmetrical manner. However, parallel deletion of simple points does not, in general, guarantee topology preservation: see for example Fig. 1a, where removing all simple points would split the object and merge two components of the background, and Fig. 1b,c,d,e where all the points are simple. In fact, such a guarantee is not obvious to obtain, even

for the 2D case (see [12], where fifteen published parallel thinning algorithms are analyzed, and counter-examples are shown for five of them). In the 2D case, a popular method due to A. Rosenfeld [31] consists of dividing each thinning step into substeps. In each substep, only simple points that have no neighbor belonging to the object in one of the four main directions (north, south, east, west) are candidate for deletion. In addition, two special configurations made of two adjacent pixels (see Fig. 1c,d) must be preserved, as well as their $\pi/2$ rotations. This strategy prevents a whole object to vanish like in Fig. 1b, or to break like in Fig. 1a. However, it cannot be straightforwardly extended to 3D. In this case, the six main directions are north, south, east, west, up and down. In Fig. 1e, the voxels x,y are simple voxels that have no neighbor belonging to the object in the direction “up”, but if we remove them in parallel, the object splits.

C. Ronse introduced the minimal non-simple sets [30] to study the conditions under which points may be removed simultaneously while preserving topology of 2D objects. This leads to verification methods for the topological soundness of parallel thinning algorithms. Such methods have been proposed for 2D algorithms by C. Ronse [30] and R. Hall [16], they have been developed for the 3D case by T.Y. Kong [22, 17] and C.M. Ma [27], as well as for the 4D case by C-J. Gau and T.Y. Kong [14, 20]. For the 3D case, one of the authors [1] introduced the notion of P-simple point as a verification method but also as a methodology to design parallel thinning algorithms [2, 11, 25, 26].

Introduced recently by one of the authors, critical kernels [3, 4] constitute a general framework in the category of abstract complexes for the study of parallel thinning in any dimension. Thanks to critical kernels, one can easily design parallel thinning algorithms that produce new types of skeletons, with specific geometrical properties, while guaranteeing their topological soundness [5, 6, 7, 8]. A new definition of a simple point is proposed in [3, 4], based on the collapse operation which is a classical tool in algebraic topology and which guarantees topology preservation.




Figure 1. (a): The object is the set of all black and gray pixels. The pixels in gray are simple, but pixels x, y for example cannot be removed in parallel while preserving topology. North pixels are marked with the letter “n”. **(b,c,d):** All pixels are simple. **(e):** All voxels are simple, the voxels x and y are both “up” voxels.

Then, the notions of an *essential face* and of a *core* of a face are used to define the *critical kernel* \mathcal{K} of a complex X . The most fundamental result proved in [3, 4] is that, if a subset Y of X contains \mathcal{K} , then X collapses onto Y , hence X and Y “have the same topology”.

In this article, we show the equivalence (up to 4D) between the notion of P-simple point and the notion of weakly crucial point, derived from the framework of critical kernels. This equivalence (Th. 22) leads to the first local characterization of P-simple points in 4D.

Furthermore, we show the equivalence (up to 4D) between the notion of MNS and the notion of crucial clique, also derived from the framework of critical kernels. This equivalence (Th. 28) leads to the first characterization of MNS which can be verified by using a polynomial method.

We also retrieve straightforwardly some previously established properties of MNSs.

In order to ease the reading, proofs of new properties are deferred to the Appendix.

1 Cubical Complexes

Abstract complexes have been promoted in particular by V. Kovalevsky [24] in order to provide a sound topological basis for image analysis. For instance, in this framework, we retrieve the main notions and results of digital topology, such as the notion of simple point.

Intuitively, a cubical complex may be thought of as a set of elements having various dimensions (e.g. cubes, squares, edges, vertices) glued together according to certain rules. In this section, we recall briefly some basic definitions on complexes, see also [5, 6, 7, 8]. We consider here n -dimensional complexes, with $0 \leq n \leq 4$.

Let S be a set. If T is a subset of S , we write $T \subseteq S$. We denote by $|S|$ the number of elements of S .

Let \mathbb{Z} be the set of integers. We consider the families of sets $\mathbb{F}_0^1, \mathbb{F}_1^1$, such that $\mathbb{F}_0^1 = \{\{a\} \mid a \in \mathbb{Z}\}$, $\mathbb{F}_1^1 = \{\{a, a+1\} \mid a \in \mathbb{Z}\}$. A subset f of \mathbb{Z}^n , $n \geq 2$, which is the Cartesian product of exactly m elements of \mathbb{F}_1^1 and $(n-m)$ elements of \mathbb{F}_0^1 is called a *face* or an m -*face* of \mathbb{Z}^n , m is the *dimension* of f , we write $\dim(f) = m$.

Observe that any non-empty intersection of faces is a face. For example, the intersection of two 2-faces A and B may be either a 2-face (if $A = B$), a 1-face, a 0-face, or the empty set.




Figure 2. Graphical representations of: (a) a 0-face, (b) a 1-face, (c) a 2-face, (d) a 3-face, (e) a 4-face.

We denote by \mathbb{F}^n the set composed of all m -faces of \mathbb{Z}^n , with $0 \leq m \leq n$. An m -face of \mathbb{Z}^n is called a *point* if $m = 0$, a *(unit) edge* if $m = 1$, a *(unit) square* if $m = 2$, a *(unit) cube* if $m = 3$, a *(unit) hypercube* if $m = 4$ (see Fig. 2).

Let f be a face in \mathbb{F}^n . We set $\hat{f} = \{g \in \mathbb{F}^n \mid g \subseteq f\}$ and $\hat{f}^* = \hat{f} \setminus \{f\}$. Any $g \in \hat{f}$ is a *face of f* , and any $g \in \hat{f}^*$ is a *proper face of f* .

If X is a finite set of faces in \mathbb{F}^n , we write $X^- = \cup\{\hat{f} \mid f \in X\}$, X^- is the *closure of X* .

A set X of faces in \mathbb{F}^n is a *cell* or an m -*cell* if there exists an m -face $f \in X$, such that $X = \hat{f}$. The *boundary of a cell \hat{f}* is the set \hat{f}^* .

A finite set X of faces in \mathbb{F}^n is a *complex (in \mathbb{F}^n)* if $X = X^-$. Any subset Y of a complex X , which is also a complex, is a *subcomplex of X* . If Y is a subcomplex of X , we write $Y \preceq X$. If X is a complex in \mathbb{F}^n , we also write $X \preceq \mathbb{F}^n$. In Fig. 3, some complexes are represented. Notice that any cell is a complex.

Let $X \subseteq \mathbb{F}^d$ be a set of faces. A sequence $\pi = \langle f_0, \dots, f_\ell \rangle$ of faces of X is a *path in X (from f_0 to f_ℓ)* if either f_i is included in f_{i+1} or f_{i+1} is included in f_i , for each $i \in \{0, \dots, \ell-1\}$.

Let $X \subseteq \mathbb{F}^d$. We say that X is *connected* if, for any two faces f, g in X , there is a path from f to g in X ; otherwise we say that X is *disconnected*. We say that Y is a (*connected*) *component of X* if $Y \neq \emptyset$, $Y \subseteq X$, Y is connected and if Y is maximal for these properties (i.e., we have $Z = Y$ whenever $Y \subseteq Z \subseteq X$ and Z is connected). Notice that the empty set is connected but has no connected component.

Let $X \subseteq \mathbb{F}^n$. A face $f \in X$ is a *facet of X* if there is no $g \in X$ such that $f \in \hat{g}^*$. We denote by X^+ the set composed of all facets of X .

If X is a complex, observe that in general, X^+ is not a complex, and that $[X^+]^- = X$.

Let $X \preceq \mathbb{F}^n$, $X \neq \emptyset$, the number $\dim(X) = \max\{\dim(f) \mid f \in X^+\}$ is the *dimension of X* . We say that X is an *m -complex* if $\dim(X) = m$.

We say that X is *pure* if, for each $f \in X^+$, we have $\dim(f) = \dim(X)$.

In Fig. 3, the complexes (a) and (f) are pure, while (b,c,d,e) are not.

2 Collapse and simple sets

Intuitively a subcomplex of a complex X is simple if its removal from X “does not change the topology of X ”. In this section we recall a definition of a simple subcomplex based on the operation of collapse [15], which is a discrete analogue of a continuous deformation (a homotopy).

Let X be a complex in \mathbb{F}^n and let $f \in X$. If there exists one face $g \in \hat{f}^*$ such that f is the only face of X that strictly includes g , then g is said to be *free for X* and the pair (f, g) is said to be a *free pair for X* . Notice that, if (f, g) is a free pair, then we have necessarily $f \in X^+$ and $\dim(g) = \dim(f) - 1$.

Let X be a complex, and let (f, g) be a free pair for X . The complex $X \setminus \{f, g\}$ is an *elementary collapse of X* .

Let X, Y be two complexes. We say that X *collapses onto Y* if $Y = X$ or if there exists a *collapse sequence from X to Y* , i.e., a sequence of complexes $\langle X_0, \dots, X_\ell \rangle$ such that $X_0 = X$, $X_\ell = Y$, and X_i is an elementary collapse of X_{i-1} , $i = 1, \dots, \ell$. If X collapses onto Y and Y is a complex made of a single point, we say that X is *collapsible*.

Fig. 3 illustrates a collapse sequence. Observe that, if X is a cell of any dimension, then X is collapsible. It may eas-

ily be seen that the collapse operation preserves the number of connected components.




Figure 3. (a): A pure 3-complex $X \preceq \mathbb{F}^3$, and a 3-face $f \in X^+$. (f): A complex Y which is the detachment of \hat{f} from X . (a-f): A collapse sequence from X to Y .

Let X, Y be two complexes. Let Z such that $X \cap Y \preceq Z \preceq Y$, and let $f, g \in Z \setminus X$. The pair (f, g) is a free pair for $X \cup Z$ if and only if (f, g) is a free pair for Z . Thus, by induction, we have the following property.

Proposition 1 ([3, 4]). *Let $X, Y \preceq \mathbb{F}^n$. The complex $X \cup Y$ collapses onto X if and only if Y collapses onto $X \cap Y$.*

The operation of detachment allows us to remove a subset from a complex, while guaranteeing that the result is still a complex.

Definition 2 ([3, 4]). *Let $Y \subseteq X \preceq \mathbb{F}^n$. We set $X \otimes Y = (X^+ \setminus Y^+)^-$. The set $X \otimes Y$ is a complex which is the detachment of Y from X .*

In the following, we will be more particularly interested in the case where Y is a single cell. For example in Fig. 3a, we see a complex X containing a 3-cell \hat{f} , and $X \otimes \hat{f}$ is depicted in Fig. 3f.

Let us now recall here a definition of simplicity ([3, 4], see also [9]) based on the collapse operation, which can be seen as a discrete counterpart of the one given by T.Y. Kong [18].

Definition 3 ([3, 4]). *Let $Y \subseteq X$; we say that Y is simple for X if X collapses onto $X \otimes Y$.*

The collapse sequence displayed in Fig. 3 (a-f) shows that the cell \hat{f} (and the face f) is simple for the complex depicted in (a).

The notion of attachment, as introduced by T.Y. Kong [17, 18], leads to a local characterization of simple sets, which follows easily from Prop. 1.

Let $Y \preceq X \preceq \mathbb{F}^n$. The *attachment* of Y for X is the complex defined by $Att(Y, X) = Y \cap (X \otimes Y)$.

Proposition 4 ([3, 4]). *Let $Y \preceq X \preceq \mathbb{F}^n$. The complex Y is simple for X if and only if Y collapses onto $Att(Y, X)$.*

Let us recall two important properties proved in [13], which will be used to establish the main results of this paper. In [13], these properties were introduced for proving new characterizations of 2D, 3D and 4D simple points.

Consider three complexes A, B, C . If A collapses onto C and A collapses onto B , then we know that A, B and C “have the same topology”. If in addition we have $C \preceq B \preceq A$, it is tempting to conjecture that B collapses onto C . Quite surprisingly this is not always true, and this fact constitutes indeed one of the principal difficulties when dealing with certain topological properties, such as the Poincaré conjecture. Classical counter-examples to this assertion are the Bing’s house and the dunce hat ([10, 32], see also [13, 29]).

In the two-dimensional discrete plane \mathbb{F}^2 , such counter-examples cannot be found, consequently the above conjecture is true in this case. We call it a confluence property. In [13] we show that, in the boundary of an n -face with $n \leq 4$, there is “not enough room” to build such counter-examples, and thus some kinds of confluence properties hold.

Theorem 5 ([13]). *Let f be a d -face with $d \in \{2, 3, 4\}$, let $A, B \preceq \hat{f}^*$ such that $B \preceq A$, and A is collapsible. Then, B is collapsible if and only if A collapses onto B .*

Theorem 6 ([13]). *Let f be a d -face with $d \in \{2, 3, 4\}$, and let $C, D \preceq \hat{f}^*$ such that $D \preceq C$, and \hat{f} collapses onto D . Then, \hat{f} collapses onto C if and only if C collapses onto D .*

It is also proved in [13] that extensions of Th. 5 and Th. 6 to dimension 5 and higher do not hold.

Th. 8 (see below) is a key property¹ which will be used to prove Prop. 19, Prop. 26 and Th. 22. Let us first state a simpler version of this property.

Proposition 7. *Let f be a d -face with $d \in \{2, 3, 4\}$, and let $X, Y \preceq \hat{f}^*$ such that X is collapsible and Y is collapsible. Then, $X \cup Y$ is collapsible if and only if $X \cap Y$ is collapsible.*

Th. 8 generalizes Prop. 7 to an arbitrary number of sub-complexes.

Theorem 8. *Let f be a d -face with $d \in \{2, 3, 4\}$, let ℓ be an integer strictly greater than 1, let X_1, \dots, X_ℓ be ℓ sub-complexes of \hat{f} . The two following assertions are equivalent:*

¹Notice that a similar property holds in \mathbb{R}^3 , in the framework of algebraic topology, if we replace the notion of collapsibility onto a point by the one of contractibility [19, 28].

i) *For all $L \subseteq \{1, \dots, \ell\}$ such that $L \neq \emptyset$, $\cup_{i \in L} X_i$ is collapsible.*

ii) *For all $L \subseteq \{1, \dots, \ell\}$ such that $L \neq \emptyset$, $\cap_{i \in L} X_i$ is collapsible.*

3 Critical kernels

Let us briefly recall the framework introduced by one of the authors (in [3, 4]) for thinning, in parallel, discrete objects with the warranty that we do not alter the topology of these objects. We focus here on the two-, three- and four-dimensional cases, in fact this method is valid for complexes of arbitrary dimension. This method is based solely on three notions, the notion of an essential face which is used to define the core of a face, and the notion of a critical face.

Definition 9 ([3, 4]). *Let $X \preceq \mathbb{F}^n$ and let $f \in X$. We say that f is an essential face for X if f is precisely the intersection of all facets of X that contain f , i.e., if $f = \cap \{g \in X^+ \mid f \subseteq g\}$. We denote by $Ess(X)$ the set composed of all essential faces of X . If f is an essential face for X , we say that \hat{f} is an essential cell for X . If $Y \preceq X$ and $Ess(Y) \subseteq Ess(X)$, then we write $Y \trianglelefteq X$.*

Observe that a facet of X is necessarily an essential face for X , i.e., $X^+ \subseteq Ess(X)$.

Definition 10 ([3, 4]). *Let $X \preceq \mathbb{F}^n$ and let $f \in Ess(X)$. The core of \hat{f} for X is the complex $Core(\hat{f}, X) = \cup \{\hat{g} \mid g \in Ess(X) \cap \hat{f}^*\}$.*

Definition 11 ([3, 4]). *Let $X \preceq \mathbb{F}^n$ and let $f \in X$. We say that f and \hat{f} are regular for X if $f \in Ess(X)$ and if \hat{f} collapses onto $Core(\hat{f}, X)$. We say that f and \hat{f} are critical for X if $f \in Ess(X)$ and if f is not regular for X .*

If $X \preceq \mathbb{F}^n$, we set $Critic(X) = \cup \{\hat{f} \mid f \text{ is critical for } X\}$, we say that $Critic(X)$ is the critical kernel of X .

A face f in X is a maximal critical face, or an M-critical face (for X), if f is a facet of $Critic(X)$.




Figure 4. A complex X for the illustration of the notions of essential face, core, critical and M-critical face (see text).

Fig. 4, where a complex X is depicted, helps us to illustrate these notions. The letters A, B, C, D, E, F indicate all the facets (2-faces), the letters a, b, c, d, e, f, g indicate some of the 1-faces, and the letters t, u, v, w, x, y, z some 0-faces. The essential faces of the complex X form the set $Ess(X) = \{A, B, C, D, E, F, a, b, c, d, t, x, y, z\}$ (essential 0-faces and 1-faces are highlighted in the figure). The core of \hat{D} for X is the set $\{x, c, y, z\}$, the core of \hat{E} for X is $\{y, d, u\}$. We can see that a collapse sequence from the cell \hat{E} to $\{y, d, u\}$ exists: e.g., $\langle \hat{E}, \{v, f, w, g, y, d, u\}, \{w, g, y, d, u\}, \{y, d, u\} \rangle$. Thus, the facet E is regular for X . On the other hand, we can see that no collapse sequence from \hat{D} to $\{x, c, y, z\}$ exists, since the latter complex is disconnected: the facet D is critical for X . We will see later that, as a consequence of Prop. 14, a facet is simple if and only if it is regular. Let us now consider the case of faces a, b and c , which are not facets. We have $Core(\hat{a}, X) = \{t\}$, $Core(\hat{b}, X) = \{t, x\}$ and $Core(\hat{c}, X) = \{x, y\}$. Thus, a is regular and b, c are critical for X . An M-critical face is a face that is critical and not included in any other critical face, thus the face b is M-critical (as it may be seen that both B and C are regular), but the face c is not M-critical, since it is included in the critical face D . See also Fig. 5a for an illustration of critical faces in 3D.

The following properties of the core of a cell will be useful in the sequel.

Proposition 12 ([3]). *Let $X \preceq \mathbb{F}^n$, and let $f \in Ess(X)$. Let $K = \{g \in X \mid f \subseteq g\}$, and let $Y = X \circ K$. We have: $Core(\hat{f}, X) = Att(\hat{f}, Y \cup \hat{f}) = \hat{f} \cap Y$.*

Corollary 13 ([3, 4]). *Let $X \preceq \mathbb{F}^n$, and let $f \in X^+$. We have: $Core(\hat{f}, X) = Att(\hat{f}, X)$.*

As shown below, there is a strong link between the notions of regular and simple face.

Proposition 14 ([3, 4]). *Let $X \preceq \mathbb{F}^n$, and let $f \in Ess(X)$. Let $K = \{g \in X \mid f \subseteq g\}$ and $Z = [X \circ K] \cup \hat{f}$. The face f is regular for X if and only if it is simple for Z .*

In particular, whenever f is a facet, f is regular for X if and only if f is simple for X (since, in this case, we have $K = \{f\}$ and thus $Z = X$).

The following theorem is the most fundamental result concerning critical kernels. We will use it in the sequel of this paper in dimension 4 or less, but notice that the theorem holds whatever the dimension.

Theorem 15 ([3, 4]). *Let $n \in \mathbb{N}$, let $X \preceq \mathbb{F}^n$.*

- i) *The complex X collapses onto its critical kernel.*
- ii) *If $Y \trianglelefteq X$ contains the critical kernel of X , then X collapses onto Y .*
- iii) *If $Y \trianglelefteq X$ contains the critical kernel of X , then any Z such that $Y \preceq Z \trianglelefteq X$ collapses onto Y .*

In Fig. 5, we show that the very notion of critical ker-




Figure 5. (a): A 3-complex X , made of 12 cubes, and its critical faces (highlighted). (b): The critical kernel $X_1 = Critic(X)$. (c): $X_2 = Critic(X_1)$. (d): $X_3 = Critic(X_2) = Critic(X_3)$.

nel can be seen as a powerful thinning algorithm, which consists of computing iteratively the critical kernel of the preceding computation. Furthermore, Th. 15ii tells us that any subcomplex $Y \trianglelefteq X$ that is “between” X (Fig. 5a) and X_1 (Fig. 5b) is such that X collapses onto Y . This true, in particular, of any subcomplex Y that is a pure 3-complex containing X_1 . This property gives birth to a wide class of parallel thinning algorithms, where different criterions, based e.g. on geometrical notions, can be used in order to choose a particular set as the result of a single thinning step (see Sec. 4 and [5, 6, 7, 8]).

4 Crucial kernels

In the image processing literature, a digital image is often considered as a set of pixels in 2D, voxels in 3D, or 4-xels in 4D. A pixel (resp. voxel, 4-xel) is an elementary square (resp. cube, hypercube), thus an easy correspondence can be made between this classical view and the framework of cubical complexes. If a complex $X \preceq \mathbb{F}^d$ is a union of d -cells, i.e., a pure d -complex, we write $X \sqsubseteq \mathbb{F}^d$. If $X, Y \sqsubseteq \mathbb{F}^d$ and $Y \preceq X$, then we write $Y \sqsubseteq X$. From now on, we consider only complexes that are unions of d -cells.

Notice that, if $X \sqsubseteq \mathbb{F}^d$ and if \hat{f} is a d -cell of X , then $X \circ \hat{f} \sqsubseteq \mathbb{F}^d$. There is indeed an equivalence between the operation on complexes that consists of removing (by detachment) a simple d -cell, and the removal of a 8-simple (resp. 26-simple, 80-simple) point in the framework of 2D (resp. 3D, 4D) digital topology (see [17, 18, 8, 6]).

If X is a pure d -complex (e.g., a union of voxels in \mathbb{F}^3), the critical kernel of X is not necessarily a pure d -complex (see Fig. 5b). The notion of crucial clique introduced in [8],

allow us to recover a pure d -subcomplex Y of an arbitrary pure d -complex X , under the constraint that X collapses onto Y .

Definition 16 ([8]). *Let $X \subseteq \mathbb{F}^d$, and let f be an M -critical face for X . The set K of all the facets of X that contain f is called a crucial clique (for X). More precisely, K is the crucial clique induced by f .*




Figure 6. Crucial cliques in \mathbb{F}^3 (represented in light gray): (a) induced by an M -critical 0-face; (b,c) induced by an M -critical 1-face; (d,e,f) induced by an M -critical 2-face. The considered M -critical faces are in dark gray, the core of these M -critical faces (when non-empty) is represented in black.

Some 3D crucial cliques are illustrated in Fig. 6. Observe that Fig. 6e depicts precisely the configuration of voxels x,y in Fig. 1e. It may be easily seen that, informally speaking, a thinning step that preserves all non-simple voxels and at least one voxel in each crucial clique, preserves topology.

The following parallel thinning scheme takes as input an “object” X that is a pure d -complex, and a subset K of X^+ , called *constraint set*, composed of facets of X that must be preserved during the thinning.

Scheme 1: CrucialThinning

Data : $d \in \{2, 3, 4\}, X \subseteq \mathbb{F}^d, K \subseteq X^+$
Result : X

- 1 **repeat**
- 2 | $D :=$ set of facets of X that are critical for X or that are in K ;
- 3 | $T :=$ set of facets of X that belong to a crucial clique included in $X^+ \setminus D$;
- 4 | $X := [D \cup T]^-$;
- 5 **until** *stability* ;

For each single step of Scheme 1, it may easily be seen that any critical face of X , if not contained in K or in a critical facet of X , is contained in T . Thus, by Th. 15, we deduce that the set X at step i collapses onto the set X at step $i + 1$, and that Scheme 1 preserves topology.

Scheme 1 is very general, since any set K may be used to constrain the thinning. Below, we give three examples of thinning methods based on this scheme, which illustrate the versatility of the critical kernels framework. Other examples may be found in [5, 6, 7, 8].

Example 1: symmetrical thinning without constraint set.

In Fig. 7 and Fig. 8, we illustrate the steps of Scheme 1 with no constraint ($K = \emptyset$). The original objects are the same as in Fig. 1a and Fig. 1e respectively. In Fig. 7, three steps are needed to reach stability, and in Fig. 8 only two steps are necessary.




Figure 7. Illustration of Scheme 1 in 2D, $K = \emptyset$.

Such minimal skeletons may be used in some applications where we are not interested to keep the branches of a skeleton while we want to keep the topology of an object and to be invariant by isometries of \mathbb{Z}^n .

Example 2: symmetrical thinning with medial axis.

A ball B is maximal for an object X if B is included in X and is not strictly included in any other ball included in X . The set of all its maximal balls is often called the medial axis of the object.




Figure 8. Illustration of Scheme 1 in 3D, $K = 0$.





Figure 9. A 2D curvilinear skeleton (top), and a 3D surfacic skeleton (bottom), obtained thanks to the CrucialThinning scheme.

In Fig. 9, we show an example of curvilinear skeleton in 2D and surfacic skeleton in 3D, obtained thanks to the above parallel thinning scheme. In both cases, the constraint set K was a set of centers of maximal balls based on the so-called Manhattan distance, more precisely those maximal balls having a radius greater than a chosen threshold (0 for the 2D object, 3 for the 3D one).

Example 3: thinning with directional strategy.

We have seen in the introduction that the directional strategy for parallel thinning, proposed by Rosenfeld in 2D [31], cannot be straightforwardly extended to the 3D case. In fact, the question of knowing whether this strategy has a “natural extension” to 3D was cited among three open questions relative to digital topology by Kong, Litherland and Rosenfeld in [23] (question 547). We show here that the critical kernels framework indeed offers a direct way to extend the directional strategy to dimensions higher than 2.

A slight variant of Scheme 1 consists of computing the set K dynamically, at each iteration, from the current state of the set X .

To implement in 3D the directional method, let us take as constraint K the set of all points that are not “up” voxels, letting as candidates for deletion only those voxels that have no neighbor belonging to the object in the “up” direction: see the result in Fig. 10b. Repeating this procedure with down, north, south, east, and west voxels achieves one step of directional thinning, as illustrated in Fig. 10. Notice that no “special configuration” (like the ones of Fig. 1c,d) need to be introduced, in 2D as in 3D. Such configurations are indeed crucial cliques, thus Scheme 1 automatically preserves them. Notice also that similar extensions may be done in dimensions higher than 3.

By way of such constraint sets, some geometric conditions such as curve end or surface border detection may also be introduced [5, 6, 7, 8], both with the symmetrical and

with the directional strategy. Let us mention that for the 3D case, a more powerful thinning scheme has been proposed in [6, 7], based on a variant of the notion of crucial clique.

It should be noted that in 2D and 3D, the methodology based on crucial cliques does not need to handle the structure of abstract complexes. In fact, we showed in [5, 8] that 2D crucial cliques may be characterized through a set of patterns (see Fig. 11) defined in the classical square grid, as in most papers related to parallel thinning. It may be seen that patterns C_1 , C_2 , C_3 and C_4 characterize crucial cliques that are connected components of the object (see Prop. 20 and Prop. 29), while C characterizes crucial cliques that are not connected components (see Prop. 30).

For example, let us consider the pixels x,y of Fig. 1a. They are both simple pixels, and they match pattern C shown in Fig. 11. Thus, we conclude that they constitute




Figure 10. Directional thinning. Voxels marked u, d, n, s are respectively up, down, north and south voxels.

a crucial clique. Indeed, it may be seen that the intersection of these two pixels (or 2-faces) is a 1-face which is critical since its core is made of two points. Furthermore this 1-face is M -critical since both x and y are simple.

Furthermore, we showed in [5, 6, 7, 8] that only one pattern in 2D (the pattern C of Fig. 11) and three patterns in 3D are sufficient, in addition to simple point detection, to implement this methodology.

Now, let us state two properties of crucial cliques which are essential for the proof of one of our main results (Th. 28).

Proposition 17. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let f be an M -critical face of X , let K be the crucial clique induced by f , and let k be any facet of K . Let K' be such that $K' \subseteq K \setminus \{k\}$ and $K' \neq K \setminus \{k\}$.*

Then, k is a simple face of the complex $X \otimes K'$.

Proposition 18. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let f be an M -critical face of X , let K be the crucial clique induced by f , and let k be any facet of K .*

Then, k is not a simple face of the complex $[X \otimes K] \cup \hat{k}$.

Prop. 19, below, plays a role in the proofs of Prop. 26 (hence also Th. 28) and Th. 20.

Proposition 19. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let $K \subseteq X^+$, $k \in K$, such that k is not simple for $[X \otimes K] \cup \hat{k}$ and k is simple for $[X \otimes K']$ whenever $K' \subseteq K \setminus \{k\}$ and $K' \neq K \setminus \{k\}$. Then, $\cap\{h \in K\}$ is a face.*

Some crucial cliques have a particularly simple structure:




Figure 11. Patterns and masks for crucial cliques. The 11 masks corresponding to these 5 patterns are obtained from them by applying any series of $\pi/2$ rotations. The label 0 indicates pixels that must belong to the background. The label P indicates pixels that must be simple pixels of the object. For mask C , at least one of the pixels marked A and at least one of the pixels marked B must be in the object.

those which are reduced to a small connected component, where each element is adjacent to each other (like those in Fig. 6a,b,d). We show that those are precisely the crucial cliques that are induced by an M -critical face having an empty core.

Theorem 20. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let $K \subseteq X^+$ be a crucial clique for X induced by an M -critical face f . Then, K^- is a connected component of X if and only if $\text{Core}(\hat{f}, X) = \emptyset$.*

5 P-simple points

In the framework of digital topology, one of the authors introduced the notion of P-simple point [2]. P-simple points can be used as a verification method, but also as a methodology to design parallel thinning algorithms [2, 11, 25, 26].

We show in this section that there exists an equivalence between the notion of P-simple points and a notion derived from the one of crucial clique. This equivalence has been stated for the 2D case in [8], here we show that it extends up to 4D.

First, we recall the definition of P-simple points. In this definition, the set C is a set of points that are “candidates for deletion”.

Let $X \sqsubseteq \mathbb{F}^n$, and let $C \subseteq X^+$. A facet $k \in C$ is said to be *P-simple* for $\langle X, C \rangle$ if k is simple for all complexes $X \otimes T$, such that $T \subseteq C \setminus \{k\}$.

For example, consider Fig. 1a and Fig. 1b, and assume in each case that C is the set of all simple facets (in gray). Then, no facet in Fig. 1b is P-simple, and all simple facets in Fig. 1a, except the ones that are delineated in bold, are P-simple for $\langle X, C \rangle$.

The following parallel thinning scheme based on P-simple points takes as input, as CrucialThinning (see Sec. 4), an “object” X which is a pure d -complex, and a

subset K of X^+ called constraint set, composed of facets of X that must be preserved during the thinning.

Scheme 2: PSimpleThinning

Data : $d \in \{2, 3, 4\}, X \sqsubseteq \mathbb{F}^d, K \subseteq X^+$
Result : X

1 **repeat**
2 $U :=$ set of facets of X that are simple for X ;
3 $V :=$ set of facets of X that are P-simple for
 $\langle X, U \setminus K \rangle$;
4 $X := X \odot V$;
5 **until** *stability* ;

For the 3D case, a local characterization that leads to a linear algorithm for testing P-simplicity has been proved in [2]. Until now, such a characterization was not available in 4D.

Definition 21 ([8]). *Let $X \sqsubseteq \mathbb{F}^n$, and let C be a set of facets of X , we set $D = X \odot C$. Let $k \in C$, we say that k is weakly crucial for $\langle X, D \rangle$ if k contains a face f that is critical for X , and such that all the facets of X containing f are in C .*

The following equivalence leads, in particular, to a local characterization and to the first algorithm for testing P-simplicity in 4D.

Theorem 22. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let C be a set of facets of X , let $D = X \odot C$. Let $k \in C$, the facet k is P-simple for $\langle X, C \rangle$ if and only if k is not weakly crucial for $\langle X, D \rangle$.*

Thus, any facet that is P-simple for $\langle X, C \rangle$ is not crucial for $\langle X, D \rangle$. The converse is not true. For example in \mathbb{F}^2 , if we consider a set X that consists of a two-pixels width ribbon (see Fig. 1b), and if we set $C = X^+$, it may be seen that the four pixels at the extremities of the ribbon are not crucial for $\langle X, D \rangle$, but also not P-simple for $\langle X, C \rangle$.

Indeed, it is possible to remove more facets with a thinning scheme that deletes simple facets that are not crucial than with one that deletes P-simple facets. In other words, the scheme CrucialThinning is more powerful than PSimpleThinning. To illustrate this, consider again the example of Fig. 1b, with no constraint set ($K = \emptyset$). One step of CrucialThinning deletes four pixels, and three steps reduces the ribbon to just two pixels, while PSimpleThinning leaves this object unchanged.

Let $Y \sqsubseteq X \sqsubseteq \mathbb{F}^n$. We say that Y is a *crucial retraction* of X if Y contains all the critical facets of X , and at least one facet of each crucial clique for X . By Th. 15, if Y is a crucial retraction of X then X collapses onto Y . In fact, despite the appearance, it is possible to check only with the notion of P-simple points whether the result of one step of a given algorithm is a crucial retraction or not. Since every critical face is included in an M-critical face, by Th. 22, we

have the following.

Proposition 23. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let T be a set of facets of X . The complex T^- is a crucial retraction of X if and only if each facet of $X^+ \setminus T$ is P-simple for $\langle X, X^+ \setminus T \rangle$.*

6 Minimal non-simple sets

In the preceding section, we saw that critical kernels, which are settled in the framework of abstract complexes, allow us to retrieve the notion of P-simple point proposed in the context of digital topology. Now, we show that the notion of minimal non-simple set can also be retrieved in the framework of critical kernels.

C. Ronse introduced in [30] the minimal non-simple sets (MNS) to propose some conditions under which simple points can be removed in parallel while preserving topology. This leads to verification methods for the topological soundness of 2D thinning algorithms [30, 16], 3D thinning algorithms [22, 17, 27], the 4D case has even been considered in [14, 19, 20].

The main result of this section (Th. 28) proves the equivalence between MNS and crucial cliques in dimensions 2, 3 and 4. This equivalence leads to the first characterization of MNS which can be verified using a polynomial method. In contrast, the very definition of a MNS (see below), as well as the characterization of Th. 24, involves the examination of all subsets of a given candidate set, e.g., a subset of a $2 \times 2 \times 2 \times 2$ block in 4D.

Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$. A sequence $\langle k_0, \dots, k_\ell \rangle$ of facets of X is said to be a *simple sequence* for X if k_0 is simple for X , and if, for any $i \in \{1, \dots, \ell\}$, k_i is simple for $X \odot \{k_j \mid 0 \leq j < i\}$. Let K be a set of facets of X , the set K is said to be *F-simple* (where ‘‘F’’ stands for facet) for X if K is empty, or if the elements of K can be ordered as a simple sequence for X . The set K is *minimal non-simple* for X if it is not F-simple for X and if all its proper subsets are F-simple. The following characterization will be used in the sequel.

Theorem 24 (adapted from Gau and Kong [14], theorem 3). *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, and let $K \subseteq X^+$. Then K is a minimal non-simple set for X if and only if the two following conditions hold:*

- i) *Each k of K is non-simple for $[X \odot K] \cup \hat{k}$.*
- ii) *Each k of K is simple for $[X \odot K']$ whenever $K' \subseteq K \setminus \{k\}$ and $K' \neq K \setminus \{k\}$.*

For example, it may be seen that the set $\{x, y\}$ in Fig. 1a, the set $\{x, y\}$ in Fig. 1e, as well as the sets displayed in Fig. 6 in light gray, are indeed minimal non-simple sets.

From Th. 24 and Prop. 19, we can directly retrieve a result previously established by Gau and Kong.

Theorem 25 (adapted from Gau and Kong [14], theorem 5). *Let $X \sqsubseteq \mathbb{F}^4$ (resp. $\mathbb{F}^3, \mathbb{F}^2$) and let $K \subseteq X^+$. If K is a minimal non-simple set for X , then K is a non-empty subset of some $2 \times 2 \times 2 \times 2$ block of sixteen 4-faces (resp. $2 \times 2 \times 2$ block of eight 3-faces, 2×2 block of four 2-faces).*

Prop. 26 and Prop. 27 are steps for the proof of the main result of this section, Th. 28.

Proposition 26. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let K be a minimal non-simple set for X , and let f be the intersection of all the elements of K . Then, f is a crucial face for X .*

Proposition 27. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let K be a minimal non-simple set for X , and let f be the intersection of all the elements of K . Then, f is an M -critical face for X and K is the crucial clique induced by f .*

If K is a crucial clique for X , then from Th. 24, Prop. 17 and Prop. 18, K is a minimal non-simple set for X . Conversely, if K is a minimal non-simple set for X , then by Prop. 27, K is a crucial clique. Thus, we have the following theorem.

Theorem 28. *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, and let $K \subseteq X^+$. Then K is a minimal non-simple set for X if and only if it is a crucial clique for X .*

We retrieve thanks to Th. 20, and by inspection of all possible configurations, a previously established property about MNSs that are connected components [30, 16, 17, 19], its most general formulation being found in [19].

Proposition 29 (adapted from Kong [19], theorem 3-2). *Let $X \sqsubseteq \mathbb{F}^d$, with $d \in \{2, 3, 4\}$, let $K \subseteq X^+$ be a connected component of X which is such that the intersection of all its d -faces is non-empty. Then, K is a minimal non-simple set for X .*

We also retrieve thanks to Th. 20, and from the observation that a face with a non-empty core has necessarily a dimension that is at least 1, the following property.

Proposition 30 (adapted from Gau and Kong [14], theorem 8-2). *Let $X \sqsubseteq \mathbb{F}^4$ (resp. $\mathbb{F}^3, \mathbb{F}^2$), let $K \subseteq X^+$ be a MNS which is not a connected component of X . Then, K is a non-empty subset of a $2 \times 2 \times 2$ block of eight 4-faces (resp. a 2×2 block of four 3-faces, a set of two 2-faces having an intersection that is a 1-face).*

Conclusion

Critical kernels constitute a powerful framework to study parallel homotopic thinning in any dimension. Indeed, the very notion of critical kernel may be seen as thinning scheme, which consists of iteratively computing the critical kernel of the result of the previous step. Critical kernels

may also be used to design new algorithms, as well as to check the topological validity of existing ones.

We demonstrated in this article that the main concepts previously introduced in order to study topology-preserving parallel thinning in the framework of digital topology, namely P-simple points and minimal non-simple sets, may be retrieved, better understood and enriched in the framework of critical kernels.

More precisely, P-simple points and minimal simple sets are particular cases in the critical kernels framework, which thus appear to constitute a unifying framework which encompasses previous works on parallel thinning.

Furthermore, in contrast with minimal non-simple sets, critical kernels provide a methodology to produce thinning algorithms which preserve topology “by construction”, and we showed in this paper that these algorithms are more powerful than those which may be designed on the basis of P-simple points.

Appendix

Proof of Prop. 7. Suppose that $X \cup Y$ is collapsible. By applying Th. 5 with $A = X \cup Y$ and $B = X$, we deduce that $X \cup Y$ collapses onto X . By Prop. 1, we deduce that Y collapses onto $X \cap Y$. And by applying Th. 5 with $A = Y$ and $B = X \cap Y$, we deduce that $X \cap Y$ is collapsible.

Now, suppose that $X \cap Y$ is collapsible. By applying Th. 5 with $A = X$ and $B = X \cap Y$, we deduce that X collapses onto $X \cap Y$. By Prop. 1, we deduce that $X \cup Y$ collapses onto Y . And since Y is collapsible, by transitivity $X \cup Y$ is collapsible. \square

Proof of Th. 8. By Prop. 7, the property holds whenever $\ell = 2$. Suppose that it holds until $\ell - 1 \geq 2$, and let us prove it for ℓ .

i \Rightarrow ii. For any $j \in \{1, \dots, \ell\}$, let $Z_j = \bigcap_{i=1}^{i=j} X_i$. For any $L \subseteq \{1, \dots, \ell\}$, $0 < |L| < \ell$, the recurrence hypothesis implies that $\bigcap_{i \in L} X_i$ is collapsible, in particular $Z_{\ell-1}$ is collapsible. It remains to prove that Z_ℓ is collapsible. For any $j \in \{1, \dots, \ell - 1\}$, let $Y_j = X_j \cup X_\ell$. For all $L \subseteq \{1, \dots, \ell - 1\}$, $L \neq \emptyset$, we see that $\bigcup_{i \in L} Y_i$ is collapsible. From the recurrence hypothesis, we deduce that $\bigcap_{i=1}^{i=\ell-1} Y_i$ is collapsible. But, $\bigcap_{i=1}^{i=\ell-1} Y_i = Z_{\ell-1} \cup X_\ell$. Since $Z_{\ell-1} \cup X_\ell$, $Z_{\ell-1}$ and X_ℓ all are collapsible, from Prop. 7 we deduce that $Z_{\ell-1} \cap X_\ell = Z_\ell$ is collapsible.

ii \Rightarrow i. The proof is obtained by exchanging intersection and union in the previous reasoning. \square

Proof of Prop. 17. Let $Z = [X \odot K']$ and $Y = [X \odot K'] \odot \hat{k}$ (see an illustration in Fig. 12). By definition of a crucial clique, the face f is included in any facet of K , thus f is contained in both Z and Y . Suppose that g is also an M -critical face for X which is in $[K']^-$, and let K'' be the cru-




Figure 12. Illustration (in \mathbb{F}^3) for the proof of Prop. 17.

cial clique for X induced by g . If $K'' \not\subseteq [K' \cup \{k\}]$, then by definition of Y , g must be included in a facet which is in Y . Otherwise $K'' \subseteq K$, hence $f \subseteq g$, and by definition of an M-critical face, $f = g$. From this we see that Y contains the critical kernel of X . Furthermore, we have $Z \triangleleft X$ (since $Z^+ \subseteq X^+ \subseteq \text{Ess}(X)$) and $Y \triangleleft X$ (idem). From Th. 15iii, we deduce that Z collapses onto Y , in other words, k is simple for Z . \square

Proof of Prop. 18. Let $W = [X \otimes K]$, $Y = W \cup \hat{k}$, and $Z = W \cup \hat{f}$ (see an illustration in Fig. 13). We have $Y \triangleleft X$




Figure 13. Illustration (in \mathbb{F}^3) for the proofs of Prop. 18 and Prop. 26.

(since $Y^+ \subseteq X^+ \subseteq \text{Ess}(X)$) and we can see that $Z \triangleleft X$, since f is M-critical, thus essential for X . From Th. 15, we deduce that Y collapses onto Z , and by Prop. 1, \hat{k} collapses onto $\hat{k} \cap Z$ (i).

Suppose that k is simple for Y , that is, Y collapses onto $Y \otimes \hat{k} = W$, thus by Prop. 1, \hat{k} collapses onto $\hat{k} \cap W$ (ii). By definition of Z and W we have $\hat{k} \cap W \preceq \hat{k} \cap Z$. Thus from (i), (ii) and Th. 6, we deduce that $\hat{k} \cap Z$ collapses onto $\hat{k} \cap W$. Remark that $\hat{k} \cap Z = [\hat{k} \cap W] \cup \hat{f}$, thus by Prop. 1, \hat{f} collapses onto $\hat{f} \cap [\hat{k} \cap W] = \hat{f} \cap W$. Again by Prop. 1, we deduce that $Z = W \cup \hat{f}$ collapses onto W , i.e., f is simple for Z , a contradiction with Prop. 14 and the fact that f is critical for X . \square

Proof of Prop. 19. Notice that the property trivially holds when $|K| = 1$, suppose now that $|K| = \ell > 1$. Let us write $K = \{k = k_1, k_2, \dots, k_\ell\}$. By hypothesis, k is simple for $\cup_{i \in L} \{[X \otimes K] \cup \hat{k} \cup \hat{k}_i\}$, $\forall L \subseteq \{2, \dots, \ell\}$ such that $L \neq \emptyset$. By Prop. 4, \hat{k} collapses onto $\cup_{i \in L} \{[X \otimes K] \cup \hat{k}_i\} \cap \hat{k} = \cup_{i \in L} \{([X \otimes K] \cup \hat{k}_i) \cap \hat{k}\}$, $\forall L \subseteq \{2, \dots, \ell\}$ such that $L \neq \emptyset$. By Th. 8 and Th. 5 (with $A = \hat{k}$), \hat{k} collapses onto $\cap_{i=2}^{\ell} \{([X \otimes K] \cup \hat{k}_i) \cap \hat{k}\} = ([X \otimes K] \cap \hat{k}) \cup (\cap_{i=2}^{\ell} \{\hat{k}_i \cap \hat{k}\}) =$

$([X \otimes K] \cap \hat{k}) \cup \hat{f}$, where $f = \cap \{h \in K\}$. Thus f cannot be empty, otherwise \hat{k} would collapse onto $[X \otimes K] \cap \hat{k}$, a contradiction with Prop. 4 and the fact that k is not simple for $[X \otimes K] \cup \hat{k}$. Since f is a non-empty intersection of faces, f is a face. \square

Proof of Th. 20. Let us denote by $|C(Z)|$ the number of connected components of any set of faces Z . The property is straightforward when $|K| = 1$, suppose now that $|K| > 1$. Let $Y = X \otimes K$. Suppose that K^- is a connected component of X , thus $K^- \cap Y = \emptyset$. Since $\hat{f} \subseteq K^-$, we have $\hat{f} \cap Y = \emptyset$. By Prop. 12, $\hat{f} \cap Y = \text{Core}(\hat{f}, X)$, hence $\text{Core}(\hat{f}, X) = \emptyset$. Conversely, suppose that $\text{Core}(\hat{f}, X) = \emptyset$, hence $\hat{f} \cap Y = \emptyset$ and $|C(Y \cup \hat{f})| = |C(Y)| + 1$. Since f is M-critical for X , the critical kernel of X is contained in $Y \cup \hat{f}$, furthermore $Y \cup \hat{f} \triangleleft X$. By Th. 15ii, X collapses onto $Y \cup \hat{f}$, hence $|C(Y \cup \hat{f})| = |C(X)| = |C(Y)| + 1$, and by definition of Y , K^- is a connected component of X . \square

The proof of Th. 22 will be given after the one of Prop. 26 for a more comfortable reading, as the two proofs share a common argument.

Proof of Prop. 26. From Th. 24 and Prop. 19, we deduce that f is a face of X . Let $\ell = |K|$, if $\ell = 1$ the property follows straightforwardly from the definitions. From now, suppose that $\ell \geq 2$. Let us write $K = \{k = k_1, k_2, \dots, k_\ell\}$, $W = [X \otimes K]$, $Y = W \cup \hat{k}$, and $Z = W \cup \hat{f}$ (see Fig. 13 for an illustration). Let L be any non-empty subset of $\{2, \dots, \ell\}$, and let us write $Y_L = W \cup \{k_i \mid i \in L\}^-$. From Th. 24ii, k is simple for $Y_L \cup \hat{k}$, that is, $Y_L \cup \hat{k}$ collapses onto Y_L . From Prop. 1, \hat{k} collapses onto $\hat{k} \cap Y_L = \hat{k} \cap [W \cup \{k_i \mid i \in L\}]^- = \hat{k} \cap [\cup \{W \cup \hat{k}_i \mid i \in L\}] = \cup \{[W \cup \hat{k}_i] \cap \hat{k} \mid i \in L\}$. From Th. 8 and Th. 5 (with $A = \hat{k}$), we deduce that \hat{k} collapses onto $\cap \{[W \cup \hat{k}_i] \cap \hat{k} \mid 2 \leq i \leq \ell\} = \hat{k} \cap [\cap \{W \cup \hat{k}_i \mid 1 \leq i \leq \ell\}] = \hat{k} \cap Z$; and by Prop. 1, $Y = \hat{k} \cup Z$ collapses onto Z . Suppose that f is not a critical face for X , i.e., \hat{f} collapses onto $\text{Core}(\hat{f}, X)$. From Prop. 12, we have $\text{Core}(\hat{f}, X) = \hat{f} \cap W$. Thus, since Y collapses onto $W \cup \hat{f}$ and \hat{f} collapses onto $\hat{f} \cap W$ (hence by Prop. 1 $W \cup \hat{f}$ collapses onto W), we deduce that Y collapses onto W , which means that k is simple for Y , a contradiction with Th. 24i. \square

Proof of Th. 22. Suppose that k is P-simple for $\langle X, C \rangle$. Let f be any face in k such that $f \neq k$ and such that the set K of all facets of X containing f is included in C . Let us write $K = \{k = k_1, k_2, \dots, k_\ell\}$. By definition of a P-simple facet, we know that k is simple for $[X \otimes K] \cup \hat{k}$, thus by Prop. 4, \hat{k} collapses onto $[X \otimes K] \cap \hat{k}$ (1). We also know that k is simple for $\cup_{i \in L} \{[X \otimes K] \cup \hat{k} \cup \hat{k}_i\}$, $\forall L \subseteq \{2, \dots, \ell\}$ such that $L \neq \emptyset$. By Prop. 4, \hat{k} collapses onto $\cup_{i \in L} \{[X \otimes K] \cup \hat{k}_i\} \cap \hat{k}$, $\forall L \subseteq \{2, \dots, \ell\}$ such that $L \neq \emptyset$. By Th. 8 and Th. 5, \hat{k} collapses onto $\cap_{i=2}^{\ell} \{[X \otimes K] \cup \hat{k}_i\} \cap \hat{k} = (\hat{k} \cap [X \otimes K]) \cup (\cap_{i=2}^{\ell} \{\hat{k}_i \cap \hat{k}\}) = ([X \otimes K] \cap \hat{k}) \cup \hat{f}$ (2). Furthermore, \hat{f} is collapsible (3) since f is a cell. From (1), (2), (3), Prop. 7 and Th. 5, we deduce that \hat{k} collapses onto $([X \otimes K] \cap \hat{k}) \cap \hat{f} = [X \otimes K] \cap \hat{f}$,

hence $[X \otimes K] \cap \hat{f}$ is collapsible (by Th. 5), and \hat{f} collapses onto $[X \otimes K] \cap \hat{f}$ (again by Th. 5). Thus by Prop. 14, f is not critical for X . Since this holds for any such face f , we conclude that k is not weakly crucial for $\langle X, D \rangle$.

Suppose now that k is not P-simple for $\langle X, C \rangle$. Thus, there exists a set $T \subseteq C \setminus \{k\}$ such that k is not simple for $X \otimes T$. Without loss of generality, we suppose that T is minimal for this property, i.e., k is simple for $X \otimes T'$ whatever the set T' strictly included in T . Let f denote the intersection of all the facets of $T \cup \{k\}$. From Prop. 19, we deduce that f is a face of X . From now on, the proof is essentially the same as the proof of Prop. 26, showing that f is a critical face, thus k is weakly crucial for $\langle X, D \rangle$. \square

Proof of Prop. 27. See Fig. 14 for an illustration. From Prop. 26 we know that f is critical for X , suppose that there exists a critical face f' for X which strictly contains f . We may assume that f' is M-critical for X (otherwise there would exist an M-critical face containing both f' and f , and this face should be chosen). Let $K' = \{k \in X \mid f' \subseteq k\}$ (by construction we have $K' \subseteq K$), $Y = [X \otimes K']$, and $Y' = Y \cup \hat{f}'$. We have $K' \neq K$, otherwise we would have $f' = f$. Let $k \in K'$ and let $Y'' = Y \cup \hat{k}$. From Th. 24ii we deduce




Figure 14. Illustration (in \mathbb{R}^3) for the proof of Prop. 27.

that Y'' collapses onto Y . By Prop. 1, it follows that \hat{k} collapses onto $\hat{k} \cap Y$ (1). The complex Y' contains, by construction, the critical kernel of X . Furthermore, since it can be seen that both k and f' are essential for X , we have $Y'' \trianglelefteq X$ and $Y' \trianglelefteq X$. By Th. 15iii, we deduce that Y'' collapses onto Y' . By Prop. 1, it follows that \hat{k} collapses onto $\hat{k} \cap Y'$ (2). From (1), (2) and Th. 6, we deduce that $\hat{k} \cap Y'$ collapses onto $\hat{k} \cap Y$, i.e., $(\hat{k} \cap Y) \cup \hat{f}'$ collapses onto $\hat{k} \cap Y$. By Prop. 1, \hat{f}' thus collapses onto $(\hat{k} \cap Y) \cap \hat{f}' = Y \cap \hat{f}'$ and by Prop. 12, $Y \cap \hat{f}' = \text{Core}(\hat{f}', X)$, hence a contradiction with the initial assumption that f' is critical for X . \square

References

[1] G. Bertrand. On P-simple points. *Comptes Rendus de l'Académie des Sciences, Série Math.*, I(321):1077–1084, 1995.

[2] G. Bertrand. Sufficient conditions for 3D parallel thinning algorithms. In *SPIE Vision Geometry IV*, volume 2573, pages 52–60, 1995.

[3] G. Bertrand. On critical kernels. Technical Report IGM2005-05, Université de Marne-la-Vallée, 2005.

[4] G. Bertrand. On critical kernels. *Comptes Rendus de l'Académie des Sciences, Série Math.*, I(345):363–367, 2007.

[5] G. Bertrand and M. Couprie. New 2d parallel thinning algorithms based on critical kernels. In *Combinatorial Image Analysis*, volume 4040 of *Lecture Notes in Computer Science*, pages 45–59. Springer, 2006.

[6] G. Bertrand and M. Couprie. A new 3D parallel thinning scheme based on critical kernels. In *Discrete Geometry for Computer Imagery*, volume 4245 of *Lecture Notes in Computer Science*, pages 580–591. Springer, 2006.

[7] G. Bertrand and M. Couprie. Three-dimensional parallel thinning algorithms based on critical kernels. *In preparation*, 2006.

[8] G. Bertrand and M. Couprie. Two-dimensional parallel thinning algorithms based on critical kernels. *Internal Report, Université de Marne-la-Vallée*, IGM2006-02, 2006. Also submitted for publication.

[9] G. Bertrand, M. Couprie, and N. Passat. 3-D simple points and simple-equivalence. 2007. Submitted.

[10] R.H. Bing. Some aspects of the topology of 3-manifolds related to the Poincaré conjecture. *Lectures on modern mathematics*, II:93–128, 1964.

[11] J. Burguet and R. Malgouyres. Strong thinning and polyhedral approximation of the surface of a voxel object. *Discrete Applied Mathematics*, 125:93–114, 2003.

[12] M. Couprie. Note on fifteen 2d parallel thinning algorithms. *Internal Report, Université de Marne-la-Vallée*, IGM2006-01, 2006.

[13] M. Couprie and G. Bertrand. New characterizations of simple points in 2D, 3D and 4D discrete spaces. Technical Report IGM2007-07, Université de Marne-la-Vallée, 2007. Submitted.

[14] C-J. Gau and T.Y. Kong. Minimal non-simple sets in 4D binary images. *Graphical Models*, 65:112–130, 2003.

[15] P. Gublin. *Graphs, surfaces and homology*. Chapman and Hall, 1981.

- [16] R.W. Hall. Tests for connectivity preservation for parallel reduction operators. *Topology and its Applications*, 46(3):199–217, 1992.
- [17] T. Y. Kong. On topology preservation in 2-D and 3-D thinning. *International Journal on Pattern Recognition and Artificial Intelligence*, 9:813–844, 1995.
- [18] T. Y. Kong. Topology-preserving deletion of 1's from 2-, 3- and 4-dimensional binary images. In Springer, editor, *Discrete Geometry for Computer Imagery*, volume 1347 of *Lecture Notes in Computer Science*, pages 3–18, 1997.
- [19] T. Y. Kong. Minimal non-simple and minimal non-cosimple sets in binary images on cell complexes. In *Discrete Geometry for Computer Imagery*, volume 4245 of *Lecture Notes in Computer Science*, pages 169–188. Springer, 2006.
- [20] T. Y. Kong and C-J. Gau. Minimal non-simple sets in 4-dimensional binary images with (8-80)-adjacency. In *International Workshop on Combinatorial Image Analysis*, pages 318–333, 2004.
- [21] T. Y. Kong and A. Rosenfeld. Digital topology: introduction and survey. *Computer Vision, Graphics and Image Processing*, 48:357–393, 1989.
- [22] T.Y. Kong. On the problem of determining whether a parallel reduction operator for n-dimensional binary images always preserves topology. In *procs. SPIE Vision Geometry II*, volume 2060, pages 69–77, 1993.
- [23] T.Y. Kong, R. Litherland, and A. Rosenfeld. Problems in the topology of binary digital images. In J. van Mill and G.M. Reed, editors, *Open problems in topology*, pages 376–385. Elsevier, 1990.
- [24] V.A. Kovalevsky. Finite topology as applied to image analysis. *Computer Vision, Graphics and Image Processing*, 46:141–161, 1989.
- [25] C. Lohou and G. Bertrand. A 3D 12-subiteration thinning algorithm based on P-simple points. *Discrete Applied Mathematics*, 139:171–195, 2004.
- [26] C. Lohou and G. Bertrand. A 3D 6-subiteration curve thinning algorithm based on P-simple points. *Discrete Applied Mathematics*, 151:198–228, 2005.
- [27] C.M. Ma. On topology preservation in 3d thinning. *Computer Vision, Graphics and Image Processing*, 59(3):328–339, May 1994.
- [28] C.R.F. Maunder. *Algebraic topology*. Dover, 1996.
- [29] N. Passat, M. Couprie, and G. Bertrand. Minimal simple pairs in the 3-d cubic grid. Technical Report IGM2007-04, Université de Marne-la-Vallée, 2007.
- [30] C. Ronse. Minimal test patterns for connectivity preservation in parallel thinning algorithms for binary digital images. *Discrete Applied Mathematics*, 21(1):67–79, 1988.
- [31] A. Rosenfeld. A characterization of parallel thinning algorithms. *Information and Control*, 29:286–291, 1975.
- [32] E.C. Zeeman. On the dunce hat. *Topology*, 2:341–358, 1964.