

GENERALIZED DUMONT-FOATA POLYNOMIALS AND ALTERNATIVE TABLEAUX

MATTHIEU JOSUAT-VERGÈS

ABSTRACT. Dumont and Foata introduced in 1976 a three-variable symmetric refinement of Genocchi numbers, which satisfies a simple recurrence relation. A six-variable generalization with many similar properties was later considered by Dumont. They generalize a lot of known integer sequences, and their ordinary generating function can be expanded as a Jacobi continued fraction.

We give here a new combinatorial interpretation of the six-variable polynomials in terms of the alternative tableaux introduced by Viennot. A powerful tool to enumerate alternative tableaux is the so-called “matrix Ansatz”, and using this we show that our combinatorial interpretation naturally leads to a new proof of the continued fraction expansion.

1. INTRODUCTION

The unsigned Genocchi numbers $\{G_{2n}\}_{n \geq 1}$ can be defined through their generating function:

$$(1) \quad \sum_{n=1}^{\infty} G_{2n} \frac{x^{2n}}{(2n)!} = x \cdot \tan\left(\frac{x}{2}\right).$$

They are related with even Bernoulli numbers B_{2n} by $G_{2n} = 2(4^n - 1)|B_{2n}|$, and they have a wide range of combinatorial properties [4, 10, 12, 14]. In the context of previous work by Carlitz, Riordan and Stein, an extension of these integers was proposed by Dumont and Foata [5]. It is defined by the recurrence $F_1(x, y, z) = 1$ and

$$(2) \quad F_n(x, y, z) = (x + y)(x + z)F_{n-1}(x + 1, y, z) - x^2F_{n-1}(x, y, z).$$

They show that the polynomial F_n is symmetric in x , y , and z , with non-negative coefficients, and such that $F_n(1, 1, 1) = G_{2n+2}$. Another nice property is that the generating function $\sum_{n=1}^{\infty} F_n t^n$ can be expanded as a J-fraction (more precision will be given further in this introduction). Gessel and Zeng [7] showed that F_{n+1} is the n th moment of some orthogonal polynomials known as *continuous dual Hahn polynomials*, which are an important sequence in the Askey-Wilson hierarchy [8].

Partially supported by the grant ANR08-JCJC-0011.

A further generalization of Genocchi numbers with many similar properties was defined by Dumont in terms of some combinatorial objects called *escaliers* [4]. It is a sequence of six-variable polynomials $\Gamma_n(x, y, z, \bar{x}, \bar{y}, \bar{z})$, or just Γ_n for short. They can be characterized by a recurrence relation which generalizes (2), and has been obtained independently by Randrianarivony [10] and Zeng [14]. For brevity, let Γ_n^+ denote $\Gamma_n(x+1, y, z, \bar{x}+1, \bar{y}, \bar{z})$.

Definition 1.1. The generalized Dumont-Foata polynomials are defined by $\Gamma_1 = 1$ and

$$(3) \quad \Gamma_n = (x + \bar{z})(y + \bar{x})\Gamma_{n-1}^+ + (x(\bar{y} - y) + \bar{x}(z - \bar{z}) - x\bar{x})\Gamma_{n-1}.$$

Quite a lot of known integer sequences appear as specializations of Γ_n [10, 11, 14]: Genocchi numbers, median Genocchi numbers, Euler numbers, median Euler numbers, Springer numbers. This polynomial Γ_n generalizes F_n since we have $\Gamma_n(x, y, z, x, y, z) = F_n(x, y, z)$.

Dumont [4] conjectured that we have the following J-fraction for $\sum \Gamma_n t^n$:

$$(4) \quad \sum_{n=1}^{\infty} \Gamma_n t^n = \frac{t}{1 - b_0 t - \frac{\lambda_1 t^2}{1 - b_1 t - \frac{\lambda_2 t^2}{\ddots}}},$$

where the parameters b_n and λ_n are defined by:

$$(5) \quad \begin{aligned} b_n &= (x+n)(\bar{y}+n) + (y+n)(\bar{z}+n) + (z+n)(\bar{x}+n) - n(n+1), \\ \lambda_n &= n(\bar{x}+y+n-1)(\bar{y}+z+n-1)(\bar{z}+x+n-1). \end{aligned}$$

This was proved independently by Randrianarivony [10] and Zeng [14] (and of course this implies the J-fraction expansion for $\sum F_n t^n$). More precisely, Randrianarivony's method consists in the study of a *Stieltjes tableau* and Zeng's method consists in calculations of *Hankel determinants*.

The main goal of this article is to give a new combinatorial interpretation of Γ_n in terms of *alternative tableaux* [9, 13] and six statistics on them, and obtain as a consequence a new proof of the continued fraction expansion (4). Alternative tableaux were introduced by Viennot [13] in the context of a model of statistical physics called Partially Asymmetric Simple Exclusion Process (PASEP), and previous work of Corteel and Williams [2]. The ‘‘matrix Ansatz’’ first appeared in [3], as a way to obtain the stationary distribution of the PASEP. In the combinatorial context, it is a method to enumerate these alternative tableaux in terms of operators satisfying certain relations. We will also describe an analog of the matrix Ansatz to enumerate escaliers, which are the combinatorial objects used by Dumont to define Γ_n in [4].

This article is organized as follows. In Section 2 we give definitions and known facts about alternative tableaux and the matrix Ansatz. In Section 3 we prove the new combinatorial interpretation of Γ_n in terms of alternative tableaux using the recurrence (3). Section 4 contains our new proof of the continued fraction expansion (4). In Section 5, we describe the analog of the matrix Ansatz to enumerate escaliers.

2. ALTERNATIVE TABLEAUX

Throughout this article we use the French convention for Young diagrams, and Young diagrams may contain rows or columns of size 0. Any Young diagram is characterized by its upper-right boundary, which is a sequence of unit steps going left or going down. We will encode this sequence by a word in the two letters D and E , so that D corresponds to the step \rightarrow and E corresponds to the step \downarrow . For example, $DDEDE$ is the Young diagram with two rows of respective lengths 2 and 3.

Definition 2.1. Let λ be a Young diagram. An *alternative tableau* of shape λ is a filling of λ such that each cell is either empty, contains an arrow \leftarrow or an arrow \downarrow , and each arrow has a clear view to the boundary. More precisely, all cells below a \downarrow in the same column (or to the left of a \leftarrow in the same row) are empty. A column (respectively, row) of an alternative tableau is *free* if it contains no \downarrow (respectively, no \leftarrow). We denote by $\text{fr}(T)$ (respectively, $\text{fc}(T)$) the number of free rows (respectively, free columns) of T . See Figure 1 for examples. We use here the notation with arrows as introduced by Nadeau [9].




FIGURE 1. Examples of alternative tableaux.

Alternative tableaux of a given shape can be enumerated via a method called matrix Ansatz. This method appeared in the context of a model of statistical physics (the partially asymmetric simple exclusion process), where it is used to derive the stationary probabilities of any state of the process.

Proposition 2.2 (Corteel-Williams [2]). *Let $\langle W|$ be a row vector, $|V\rangle$ a column vector, and D and E matrices such that:*

$$(6) \quad \langle W|V\rangle = 1, \quad \langle W|E = \bar{x}\langle W|, \quad D|V\rangle = y|V\rangle, \quad \text{and} \quad DE - ED = D + E.$$

Let w be a word in the two letters D and E , then we have

$$(7) \quad \langle W|w|V \rangle = \sum_T \bar{x}^{\text{fr}(T)} y^{\text{fc}(T)}$$

where the sum is over alternative tableaux T of shape w .

The result of Corteel and Williams was actually stated in terms of *permutation tableaux*, which are slightly different objects. We refer to Viennot [13] and Nadeau [9] for the corresponding statement in terms of alternative tableaux, and for the bijection between permutation tableaux and alternative tableaux. We have chosen to use alternative tableaux in this work because of their symmetry. Indeed, there is an elementary involution on alternative tableaux which is *conjugation*. To conjugate a tableau, take the image of the whole picture with respect to the South-West to North-East axis symmetry (in particular, the \leftarrow and \downarrow are exchanged). See [9] for details.

Note that relations (6) ensure that $\langle W|w|V \rangle$ is well-defined and can be computed explicitly. Indeed, we can use $DE - ED = D + E$ to obtain some coefficients $c_{i,j}$ such that $w = \sum_{i,j} c_{i,j} E^i D^j$, and from the other relations we can obtain $\langle W|w|V \rangle$. We refer to [2, 1] for more details.

Although not necessary to compute $\langle W|w|V \rangle$ for a given word w , it is useful to have explicit matrices satisfying the PASEP matrix Ansatz. It can be checked [3] that the following $\mathbb{N} \times \mathbb{N}$ -matrices:

$$(8) \quad D = \begin{pmatrix} y & 1 & & & (0) \\ & y+1 & 2 & & \\ & & y+2 & 3 & \\ & & & y+3 & \ddots \\ (0) & & & & \ddots \end{pmatrix}, E = \begin{pmatrix} \bar{x} & & & & (0) \\ y+\bar{x} & \bar{x}+1 & & & \\ & y+\bar{x}+1 & \bar{x}+2 & & \\ & & y+\bar{x}+2 & \bar{x}+3 & \\ (0) & & & & \ddots & \ddots \end{pmatrix},$$

satisfy $DE - ED = D + E$. They are essentially a particular case of matrices defined by Derrida & al [3] in the context of the PASEP. As for the vectors, we can take $\langle W| = (1, 0, 0, \dots)$ and $|V \rangle = (1, 0, 0, \dots)^*$, and all relations in (6) are satisfied.

3. THE NEW COMBINATORIAL INTERPRETATION OF Γ_n

It is known [13] that G_{2n+2} is the number of alternative tableaux whose shape is the staircase with n rows and columns, *i.e.* the Young diagram corresponding to the word $(DE)^n$. In [1], we have given three statistic in staircase alternative tableaux to give a combinatorial interpretation of $F_n(x, y, z)$. These are: the number of free rows, the number of free columns, and the number of corners containing \leftarrow or \downarrow . Here, we give six statistics for the more general case of Γ_n . Another difference

is that in [1], the combinatorial interpretation was derived from the J-fraction for $\sum F_n t^n$, but here we use the recurrence relation (3) to prove the result.

Definition 3.1. A column (respectively, row) of an alternative tableau is *empty* if it contains no \downarrow nor \leftarrow . Let T be an alternative tableau. We denote by:

- $\text{emr}(T)$, the number of empty rows in T ,
- $\text{fnc}(T)$, the number of free non-empty columns in T ,
- $\text{dco}(T)$, the number of corners containing a \downarrow in T ,
- $\text{fnr}(T)$, the number of free non-empty rows in T ,
- $\text{emc}(T)$, the number of empty columns in T ,
- $\text{lco}(T)$, the number of corners containing a \leftarrow in T .

Moreover let \mathcal{T}_n be the set of alternative tableaux whose shape is the staircase Young diagram with n rows and n columns.

For example, the rightmost tableau in Figure 1 is in \mathcal{T}_5 , and the six statistics that we have just defined are respectively 2, 2, 2, 0, 1, 0. The main new result of this article is the following:

Theorem 3.2. *For any $n \geq 1$, we have*

$$(9) \quad \Gamma_n(x, y, z, \bar{x}, \bar{y}, \bar{z}) = \sum_{T \in \mathcal{T}_{n-1}} x^{\text{emr}(T)} y^{\text{fnc}(T)} z^{\text{dco}(T)} \bar{x}^{\text{fnr}(T)} \bar{y}^{\text{emc}(T)} \bar{z}^{\text{lco}(T)}.$$

Proof. Both sides are equal to 1 when $n = 1$, so it suffices to show that the right-hand side satisfies the recurrence relation (3). We distinguish six kinds of tableaux in the set \mathcal{T}_{n-1} , according to the content of their leftmost column and upper left corner. Assuming that the theorem is true for $n - 1$, we will show that these six kinds of tableaux have generating functions which add up to the right-hand side of (3). This is summarized in the following table.

The upper left corner contains:

		\downarrow	\leftarrow	nothing
The leftmost column is:	empty	\times	\times	Case 4 $x\bar{y}\Gamma_{n-1}$
	free non-empty	\times	Case 2 $y\bar{z}\Gamma_{n-1}^+$	Case 5 $xy(\Gamma_{n-1}^+ - \Gamma_{n-1})$
	non-free	Case 1 $\bar{x}z\Gamma_{n-1}$	Case 3 $\bar{x}\bar{z}(\Gamma_{n-1}^+ - \Gamma_{n-1})$	Case 6 $x\bar{x}(\Gamma_{n-1}^+ - \Gamma_{n-1})$

For example we will show that the tableaux of the fourth kind (case 4), *i.e.* those having an empty leftmost column, have generating function $x\bar{y}\Gamma_{n-1}$. The three cells containing a \times in this table do not correspond to any tableaux.




FIGURE 2. Recursive construction of staircase alternative tableaux.

- Case 1. When the upper-left corner contains a \downarrow , there is no other arrow in the leftmost column. This corresponds to the first picture in Figure 2. In this first kind of tableaux, the topmost row is free non-empty, and the upper left corner contains a \downarrow , so this gives a factor $\bar{x}z$. After removing the leftmost column, there can remain any tableau in \mathcal{T}_{n-2} , hence the factor Γ_{n-1} . So the first kind of tableaux gives indeed the term $\bar{x}z\Gamma_{n-1}$.
- Case 2. There is a factor $y\bar{z}$ since we assume that there is a \leftarrow in the upper left corner, and that the leftmost column is free and non-empty. These tableaux can be obtained the following way: consider any tableau T in \mathcal{T}_{n-2} , add a column to its left with a \leftarrow in the topmost cell of the added column. We color some of the other cells in the added column in gray as in the second picture in Figure 2, such that a cell is colored if there is no \leftarrow to its right. Then, decide whether each gray cell is empty or contains a \leftarrow . All tableaux of the second kind can be obtained this way, and the gray cells are in correspondence with the free rows of T . At the level of generating functions, this amounts to substitute x with $x+1$ and \bar{x} with $\bar{x}+1$. Indeed, an empty (respectively, free non-empty) row remains so if we add nothing in the gray cell, but becomes non-free if we add a \leftarrow .
- Case 3. This corresponds to the second picture in Figure 2, but with the assumption that there is a \downarrow in one of the gray cells. Let us consider the set S of tableaux of the second kind such that there is at least a \leftarrow in some gray cell. This set has generating function $y\bar{z}(\Gamma_{n-1}^+ - \Gamma_{n-1})$, indeed we have already $y\bar{z}\Gamma_{n-1}^+$ for the all tableaux of the second kind and the term $-y\bar{z}\Gamma_{n-1}$ removes the cases where all gray cells are empty. Then, there is a bijection between this set S and the tableaux of the third kind. Indeed, let $T \in S$, consider the bottommost \leftarrow in the leftmost column of T , and replace this \leftarrow with a \downarrow . This way, we obtain exactly the tableaux of the third kind. Replacing the \leftarrow with \downarrow gives a factor $\bar{x}y^{-1}$ at the level of generating functions. Thus we obtain $\bar{x}\bar{z}(\Gamma_{n-1}^+ - \Gamma_{n-1})$ for the third kind of tableaux.
- Case 4. This is similar to case 1, and corresponds to the third picture in Figure 2. Here, removing the first column gives a factor $x\bar{y}$ since we assume

that the leftmost column is empty, and hence the upper row is empty. There can remain any tableau in \mathcal{T}_{n-2} , hence the factor Γ_{n-1} . Thus we obtain $x\bar{y}\Gamma_{n-1}$ for the fourth kind of tableaux.

- Case 5. This corresponds to the fourth picture in Figure 2, with the assumption that the gray cells contain no \downarrow and at least a \leftarrow . Here the gray cells are obtained exactly as in case 2 above. Proceeding similar to case 3 above, we obtain $xy(\Gamma_{n-1}^+ - \Gamma_{n-1})$ for the fifth kind of tableaux.
- Case 6. There is a bijection between the fifth kind and the sixth kind of tableaux, similar to the bijection used in case 3. From a tableau of the fifth kind, consider the bottommost \leftarrow in the leftmost column, and replace it with a \downarrow . Replacing the \leftarrow with \downarrow gives a factor $\bar{x}y^{-1}$ at the level of generating functions. Thus we obtain $x\bar{x}(\Gamma_{n-1}^+ - \Gamma_{n-1})$ for the sixth kind of tableaux.

Adding the six terms in the above table, we get the right-hand side of (3). This shows that the right-hand side of (9) satisfies the same recurrence as Γ_n , and completes the proof. \square

As previously mentioned, there is a simple bijection between alternative tableaux and permutation tableaux and it is possible to derive a combinatorial interpretation of Γ_n in terms of permutation tableaux, but the result is much more natural with the alternative tableaux. In particular, the conjugation of alternative tableaux gives an easy way to prove a symmetry property of Γ_n , which has been first given by Randrianarivony [10] and Zeng [14].

Proposition 3.3 ([10, 14]). *For any permutation $\sigma = u, v, w$ of x, y, z we have:*

- *if σ has signature 1, then $\Gamma_n(u, v, w, \bar{u}, \bar{v}, \bar{w}) = \Gamma_n(x, y, z, \bar{x}, \bar{y}, \bar{z})$,*
- *if σ has signature -1, then $\Gamma_n(u, v, w, \bar{u}, \bar{v}, \bar{w}) = \Gamma_n(\bar{x}, \bar{y}, \bar{z}, x, y, z)$.*

In particular, $F_n(x, y, z) = \Gamma_n(x, y, z, x, y, z)$ is symmetric in x, y , and z .

Proof. From the recurrence relation (3), we have

$$(10) \quad \Gamma(x, y, z, \bar{x}, \bar{y}, \bar{z}) = \Gamma(\bar{x}, \bar{z}, \bar{y}, x, z, y).$$

From the combinatorial interpretation in (9) and using the conjugation of alternative tableaux, we have

$$(11) \quad \Gamma(x, y, z, \bar{x}, \bar{y}, \bar{z}) = \Gamma(\bar{y}, \bar{x}, \bar{z}, y, x, z).$$

All symmetries given in the can be obtained by combining (10) and (11). \square

It is rather curious that one symmetry is obvious on the combinatorial interpretation, and another one in the recurrence relation. In the model in terms of escaliers (see Section 5), only one symmetry is obvious, and it is both in the combinatorial interpretation and in the recurrence relation. Note that any symmetry of a generating function necessarily appear in the coefficients of its expansion as a J-fraction, and indeed it is straightforward to check that the coefficients b_n and λ_n defined in (5) have the same symmetries as Γ_n .

4. THE CONTINUED FRACTION EXPANSION

In this section, we show that Γ_n can be calculated via the matrix Ansatz, and we derive as a consequence a new proof of the continued fraction expansion for $\sum \Gamma_n t^n$. We consider the matrix

$$(12) \quad M = ED + (\bar{z} + x - \bar{x})D + (z + \bar{y} - y)E + (\bar{y} - y)(x - \bar{x})I,$$

where D, E are defined in (8), and I is the identity matrix. It turns out that we can exploit Proposition 2.2 to obtain the following:

Proposition 4.1. *For any $n \geq 0$, we have*

$$(13) \quad \Gamma_{n+1}(x, y, z, \bar{x}, \bar{y}, \bar{z}) = \langle W | M^n | V \rangle.$$

To prove this, we need a few helpful definitions and lemmas.

Definition 4.2. Let \mathcal{T}_n^* be the set of pairs (T, X) where $T \in \mathcal{T}_n$ and X is a subset of the empty rows and columns of T . Such a pair (T, X) is called an *extended tableau*, and will be represented the following way: from a picture of T , each row or column in X is distinguished by a dashed line going through it. See Figure 3 for some examples. Given $U = (T, X) \in \mathcal{T}_n^*$, we define

- $\text{hr}(U)$ as the number of dashed rows,
- $\text{hc}(U)$ as the number of dashed columns.

For any statistic “stat” on alternative tableaux, and $U = (T, X) \in \mathcal{T}_n^*$ we define $\text{stat}(U) = \text{stat}(T)$. For any extended tableau U , we define the weight $w(U)$ as

$$(14) \quad \begin{aligned} w(U) &= \bar{x}^{\text{emr}(U) - \text{hr}(U)} y^{\text{fnc}(U)} z^{\text{dco}(U)} \bar{x}^{\text{fnr}(U)} y^{\text{emc}(U) - \text{hc}(U)} \bar{z}^{\text{lco}(U)} (x - \bar{x})^{\text{hr}(U)} (\bar{y} - y)^{\text{hc}(U)} \\ &= y^{\text{fc}(U) - \text{hc}(U)} z^{\text{dco}(U)} \bar{x}^{\text{fr}(U) - \text{hr}(U)} \bar{z}^{\text{lco}(U)} (x - \bar{x})^{\text{hr}(U)} (\bar{y} - y)^{\text{hc}(U)}, \end{aligned}$$

the latest equality following from $\text{emr}(T) + \text{fnr}(T) = \text{fr}(T)$ and $\text{emc}(T) + \text{fnc}(T) = \text{fc}(T)$ for any alternative tableau T .




FIGURE 3. Example of two extended alternative tableaux.

Lemma 4.3. *We have:*

$$(15) \quad \sum_{U \in \mathcal{T}_n^*} w(U) = \sum_{T \in \mathcal{T}_n} x^{\text{emr}(T)} y^{\text{fnc}(T)} z^{\text{dco}(T)} \bar{x}^{\text{fnr}(T)} \bar{y}^{\text{emc}(T)} \bar{z}^{\text{lco}(T)}.$$

Proof. In the sum $\sum_{U \in \mathcal{T}_n^*} w(U)$, we have distinguished two kinds of empty rows (dashed or non-dashed) with respective weights $x - \bar{x}$ and \bar{x} instead of one kind of empty row with weight x . Similarly we have distinguished two kinds of empty columns (dashed or non-dashed) with respective weights $\bar{y} - y$ and y instead of one kind of empty column with weight \bar{y} . By an elementary argument, it is clear that these distinctions do not change the generating function. \square

Definition 4.4. Let (T, X) be an extended tableau in \mathcal{T}_n^* . The *profile* of (T, X) is the sequence (i_1, \dots, i_n) , where:

- $i_k = 1$ if the k th corner of T is empty,
- $i_k = 2$ if the k th corner of T contains a \leftarrow ,
- $i_k = 3$ if the k th corner of T is in a dashed row but not in a dashed column,
- $i_k = 4$ if the k th corner of T contains a \downarrow ,
- $i_k = 5$ if the k th corner of T is in a dashed column but not in a dashed row,
- $i_k = 6$ if the k th corner of T is in a dashed column and in a dashed row.

Here the corners are numbered from the upper left one to the lower right one. For example, the first extended tableau in Figure 3 has profile $(1, 5, 1, 4, 6)$, and the second one has profile $(5, 2, 3, 1, 4)$.

Lemma 4.5. Let M_1, \dots, M_6 be the matrices

$$(16) \quad \begin{aligned} M_1 &= ED, & M_2 &= \bar{z}D, & M_3 &= (x - \bar{x})D, \\ M_4 &= zE, & M_5 &= (\bar{y} - y)E, & M_6 &= (\bar{y} - y)(x - \bar{x})I. \end{aligned}$$

For any $(i_1, \dots, i_n) \in \{1, \dots, 6\}^n$, we have

$$(17) \quad \sum_U w(U) = \langle W | M_{i_1} \dots M_{i_n} | V \rangle,$$

where the sum is over extended tableaux U of profile (i_1, \dots, i_n) .

Proof. Let w be the word obtained from $i_1 \dots i_n$ through the substitution $1 \mapsto ED$, $2 \mapsto D$, $3 \mapsto D$, $4 \mapsto E$, $5 \mapsto E$, $6 \mapsto \epsilon$ (ϵ being the empty word). The main point is that there is a bijection ϕ between elements in \mathcal{T}_n^* of profile (i_1, \dots, i_n) and alternative tableaux of shape w . Indeed, to build an extended tableau, once the contents of the corners are specified it remains only to choose an alternative tableau of a smaller shape. More precisely, the bijection can be done the following way:

- for each empty corner of the extended tableau, remove the corresponding cell in the Young diagram,
- shrink each dashed row or column,
- for each corner of the extended tableau containing a \leftarrow (respectively, \downarrow), shrink the row (respectively, column) containing it.

FIGURE 4. Images of extended tableaux by the map ϕ .

See Figure 4 for an example, where we give the image of the two extended tableaux in Figure 3.

The weight of an extended tableau U of profile (i_1, \dots, i_n) is the product

$$y^a z^b \bar{x}^c \bar{z}^d (x - \bar{x})^e (\bar{y} - y)^f,$$

where:

- a is the number of non-dashed free columns in U ,
- $b = \text{dco}(U)$ is the number of 4's in (i_1, \dots, i_n) ,
- c is the number of non-dashed free rows in U ,
- $d = \text{lco}(U)$ is the number of 2's in (i_1, \dots, i_n) ,
- $e = \text{hr}(U)$ is the number of 3's plus the number of 6's in (i_1, \dots, i_n) ,
- $f = \text{hc}(U)$ is the number of 5's plus the number of 6's in (i_1, \dots, i_n) .

An important property of the bijection ϕ is that the free rows (respectively, columns) of $\phi(U)$ are in correspondence with non-dashed free rows (respectively, columns) of U . It follows that

$$(18) \quad \sum_U w(U) = z^b \bar{z}^d (x - \bar{x})^e (\bar{y} - y)^f \sum_T \bar{x}^{\text{fr}(T)} y^{\text{fc}(T)},$$

where the first sum is over extended tableau of profile (i_1, \dots, i_n) and the second one is over alternative tableau of shape w .

Now, examine the product $M_{i_1} \dots M_{i_n}$. The factors D and E in this product readily gives the word w , and the other factors readily gives $z^b \bar{z}^d (x - \bar{x})^e (\bar{y} - y)^f$, so:

$$(19) \quad \langle W | M_{i_1} \dots M_{i_n} | V \rangle = z^b \bar{z}^d (x - \bar{x})^e (\bar{y} - y)^f \langle W | w | V \rangle.$$

Using Proposition 2.2, the result follows from (18) and (19). \square

Now, we can prove Proposition 4.1.

Proof. Since $M = \sum_{i=1}^6 M_i$, the expansion of M^n is also the sum of all products $M_{i_1} \dots M_{i_n}$ where (i_1, \dots, i_n) runs through the set $\{1, \dots, 6\}^n$. Hence,

$$(20) \quad \langle W | M^n | V \rangle = \sum_{(i_1, \dots, i_n) \in \{1, \dots, 6\}^n} \langle W | M_{i_1} \dots M_{i_n} | V \rangle.$$

Using Equation (17) in Lemma 4.5, this gives

$$(21) \quad \langle W|M^n|V \rangle = \sum_{U \in \mathcal{T}_{n-1}^*} w(U).$$

Using Equation (15) in Lemma 4.3 and Theorem 3.2, this is equal to Γ_{n+1} . \square

From the definitions of D and E in (8), the matrix M defined in (12) can be calculated explicitly and we obtain the following statement.

Proposition 4.6. *The matrix $M = (M_{i,j})_{i,j \in \mathbb{N}}$ is tridiagonal, and such that for any $i \geq 0$ we have*

$$(22) \quad M_{i,i} = b_i \quad \text{and} \quad M_{i,i+1}M_{i+1,i} = \lambda_{i+1},$$

where b_i and λ_i are defined in (5).

Proof. We have

$$\begin{aligned} M_{i,i} &= E_{i,i}D_{i,i} + E_{i,i-1}D_{i-1,i} + (\bar{z} + x - \bar{x})D_{i,i} + (z + \bar{y} - y)E_{i,i} + (\bar{y} - y)(x - \bar{x}) \\ &= (\bar{x} + i)(y + i) + (y + \bar{x} + i - 1)i + (\bar{z} + x - \bar{x})(y + i) \\ &\quad + (z + \bar{y} - y)(\bar{x} + i) + (\bar{y} - y)(x - \bar{x}) \\ &= x\bar{y} + y\bar{z} + z\bar{x} + i(\bar{x} + \bar{y} + \bar{z} + x + y + z) + i(2i - 1) = b_i. \end{aligned}$$

We have also

$$\begin{aligned} M_{i,i+1} &= E_{i,i}D_{i,i+1} + (\bar{z} + x - \bar{x})D_{i,i+1} \\ &= (\bar{x} + i)(i + 1) + (\bar{z} + x - \bar{x})(i + 1) = (x + \bar{z} + i)(i + 1), \end{aligned}$$

and

$$\begin{aligned} M_{i+1,i} &= E_{i+1,i}D_{i,i} + (z + \bar{y} - y)E_{i+1,i} = (y + \bar{x} + i)(y + i) + (z + \bar{y} - y)(y + \bar{x} + i) \\ &= (\bar{x} + y + i)(z + \bar{y} + i). \end{aligned}$$

Hence, $M_{i,i+1}M_{i+1,i} = \lambda_{i+1}$. It is straightforward to check that other coefficients in M are 0, and this completes the proof. \square

As a direct consequence of Propositions 4.1 and 4.6, let us give a new proof of the continued fraction expansion given in (4). First, note that $\langle W|M^n|V \rangle$ is the top-left coefficient $(M^n)_{0,0}$ of the matrix M^n . This coefficient can be obtained by expanding the product M^n and we obtain

$$(23) \quad \langle W|M^n|V \rangle = \sum_{i_1, \dots, i_{n-1} \geq 0} M_{0,i_1}M_{i_1,i_2} \cdots M_{i_{n-2},i_{n-1}}M_{i_{n-1},0}.$$

Since the matrix M is tridiagonal, we can restrict the sum to the case where two successive indices differ by at most 1, *i.e.* $|i_j - i_{j+1}| \leq 1$ for any $j \in \{0, \dots, n-1\}$ where $i_0 = i_n = 0$. These indices thus define the successive heights of a Motzkin path. Then (23) shows that Γ_{n+1} can be seen as the generating function of Motzkin paths of n steps, with a weight b_i for a level step at height i , and a weight λ_i for

a step \nearrow between height $i - 1$ and i . By a standard argument [6] this implies the continued fraction given in (4).

5. THE MATRIX ANSATZ FOR ESCALIERS

In the previous section, we have applied the link between alternative tableaux and matrices D and E satisfying $DE - ED = D + E$ to obtain the continued fraction. In this section, we consider *escaliers*, which are the combinatorial objects used by Dumont [4] to define Γ_n . We will show that these objects can be enumerated by a similar method, but with matrices B and A satisfying $BA - AB = A + I$.

We will denote a Young diagram by a word in B and A in the same way that we did with D and E (B and A respectively correspond to steps \rightarrow and \downarrow in the North-East boundary of the Young diagram).

Definition 5.1. A *surjective pretableau* is a partial filling a Young diagram with \times , such that there is at least one \times in each row and at most one \times in each column. A *surjective tableau* is a surjective pretableau such that there is exactly one \times in each column. An *escalier* (of size n) is a surjective tableau of shape $(BBA)^n$. See Figure 5 for some examples.



FIGURE 5. Examples of surjective pretableaux of shape $(BBA)^4$.

The definition of Γ_n by Dumont [4] is given in terms of escaliers of size n and six statistics on them. There is an obvious bijection between escaliers of size n and surjective pretableau of shape $(BBA)^{n-1}$ (remove the bottom row of the escalier), so that Dumont's definition is equivalent to the following (as mentioned in the introduction, this is also known to be equivalent with Definition 1.1).

Definition 5.2. A *co-corner* of the Young diagram $(BBA)^n$ is a cell which is the left neighbor of a corner (for example the upper-left cell is a co-corner). Let \mathcal{S}_n be the set of surjective pretableau of shape $(BBA)^n$. A column is *empty* if it contains no \times . A \times is *doubled* if there is another \times in the same row. We denote by:

- $\text{mi}(T)$, the number of empty columns of odd index,
- $\text{fd}(T)$, the number of corners containing a doubled \times ,
- $\text{snd}(T)$, the number of co-corners containing a non-doubled \times ,
- $\text{mp}(T)$, the number of empty columns of even index,
- $\text{fnd}(T)$, the number of corners containing a non-doubled \times ,
- $\text{sd}(T)$, the number of co-corners containing a doubled \times ,

Eventually, Γ_n can be defined as

$$(24) \quad \Gamma_n = \sum_{T \in \mathcal{S}_{n-1}} x^{\text{mi}(T)} y^{\text{fd}(T)} z^{\text{snd}(T)} \bar{x}^{\text{mp}(T)} \bar{y}^{\text{fnd}(T)} \bar{z}^{\text{sd}(T)}.$$

For example, the values of the six statistics on the first surjective pretableau in Figure 5 are 1, 1, 1, 2, 1, and 0. As for the second one, the values are 0, 1, 2, 2, 0, and 1. The fact that these objects also follow the recurrence (3) is seen by distinguishing several kinds of elements in \mathcal{S}_{n-1} according to the content of the bottom row [10, 14].

The analog of the matrix Ansatz for escaliers is given in the following proposition. The proof is similar to the case of alternative tableaux, and various examples of this kind of results were given in [1].

Proposition 5.3. *Let $\langle W|$ be a row vector, $|V\rangle$ a column vector, and A and B matrices such that:*

$$(25) \quad \langle W|V\rangle = 1, \quad \langle W|A = 0, \quad B|V\rangle = 0, \quad BA - AB = A + I.$$

Let w be a word in the two letters B and A , then the number of surjective tableaux of shape w is $\langle W|w|V\rangle$.

Proof. This is done by a recurrence on the number of cells in the Young diagram. If there is no cell, then $w = A^i B^j$ for some i and j , so $\langle W|w|V\rangle$ equals 0 if $i > 0$ or $j > 0$ and 1 otherwise. Since there is at least a \times in each row and column of a surjective tableau, there is no such tableau of shape $A^i B^j$ if $i > 0$ or $j > 0$, but we do have the “empty” surjective tableau in the “empty Young diagram” when $i = j = 0$.

Next, consider a word w which is not in the form $A^i B^j$. It means we can factorize it in $w = w_1 B A w_2$, and the factor BA corresponds to a corner of the Young diagram. We can distinguish three kinds of surjective tableaux of shape w , depending on the content of this corner.

- If the corner is empty, we can remove it and obtain any surjective tableaux of shape $w_1 A B w_2$. By the recurrence assumption, their number is $\langle W|w_1 A B w_2|V\rangle$.
- If the corner contains a doubled \times , we can delete the corner and its column, and obtain any surjective tableau of shape $w_1 A w_2$. Their number is $\langle W|w_1 A w_2|V\rangle$.
- If the corner contains a non-doubled \times , we can remove the corner and its row and column, and obtain any surjective tableau of shape $w_1 w_2$. Their number is $\langle W|w_1 w_2|V\rangle$.

It follows that the number of surjective tableaux of shape w is

$$\langle W|w_1 A B w_2|V\rangle + \langle W|w_1 A w_2|V\rangle + \langle W|w_1 w_2|V\rangle = \langle W|w_1 B A w_2|V\rangle = \langle W|w|V\rangle,$$

and this completes the recurrence. \square

It is easily checked that the following $\mathbb{N} \times \mathbb{N}$ -matrices:

$$(26) \quad B = \begin{pmatrix} 0 & 1 & & (0) \\ & 1 & 2 & \\ & & 2 & 3 \\ & & & 3 \\ (0) & & & \ddots \end{pmatrix}, \quad A = \begin{pmatrix} 0 & & & (0) \\ 1 & 0 & & \\ & 1 & 0 & \\ & & 1 & 0 \\ (0) & & \ddots & \ddots \end{pmatrix},$$

satisfy $BA - AB = A + I$. We keep the definition of $\langle W|$ and $|V\rangle$ as in the previous sections, since this also ensures that we have $\langle W|A = 0$ and $B|V\rangle = 0$.

To see how to use this result in the case of Γ_n , let us begin with the particular case $y = z = \bar{y} = \bar{z} = 1$. We know that the number of surjective tableaux of shape $(BBA)^n$ is $\langle W|(BBA)^n|V\rangle$. If we want to count surjective pretableaux, we have to authorize empty columns, and this is done by replacing B with $B + I$. Indeed, in the expansion of the product $\langle W|((B + I)(B + I)A)^n|V\rangle$, the choice of B or I in some factor corresponds to the choice of leaving a column empty or not. So the number of surjective pretableaux of shape $(BBA)^n$ is $\langle W|((B + I)(B + I)A)^n|V\rangle$. If we want to follow the empty columns of odd (respectively, even) index by the parameter x (respectively \bar{x}), it suffices to mark the terms I , and we obtain $\Gamma_{n+1}(x, 1, 1, \bar{x}, 1, 1) = \langle W|((B + xI)(B + \bar{x}I)A)^n|V\rangle$.

As for the general case, in the same way that we have obtained Proposition 4.1 from the combinatorial interpretation in terms of alternative tableaux, we can obtain the following from the combinatorial interpretation in terms of escaliers.

Proposition 5.4. *For any $n \geq 0$, we have $\Gamma_{n+1} = \langle W|N^n|V\rangle$ where N is the matrix*

$$(27) \quad N = A(B + xI)(B + \bar{x}I) + y\bar{z}(A + I) + (zI + \bar{z}A)(B + \bar{x}I) + (\bar{y}I + yA)(B + xI).$$

As in the previous section, we need some helpful definitions and lemmas.

Definition 5.5. The *profile* of $T \in \mathcal{S}_n$ is the sequence (i_1, \dots, i_n) , where

- $i_k = 1$ if the k th co-corner and k th corner are empty,
- $i_k = 2$ if the k th co-corner and k th corner both contain a \times ,
- $i_k = 3$ if the k th co-corner contains a non-doubled \times ,
- $i_k = 4$ if the k th co-corner contains a doubled \times and the k th corner is empty,
- $i_k = 5$ if the k th corner contains a non-doubled \times ,
- $i_k = 6$ if the k th corner contains a doubled \times and the k th co-corner is empty.

For example, the two surjective pretableaux in Figure 5 have respective profile $(5, 6, 3, 1)$ and $(3, 3, 6, 4)$.

Lemma 5.6. *We define the matrices $N_1 = A(B + xI)(B + \bar{x}I)$, $N_2 = y\bar{z}(A + I)$, $N_3 = z(B + \bar{x}I)$, $N_4 = \bar{z}A(B + \bar{x}I)$, $N_5 = \bar{y}(B + xI)$, $N_6 = yA(B + xI)$. For any*

$(i_1, \dots, i_n) \in \{1, \dots, 6\}^n$, we have

$$(28) \quad \sum_T x^{\text{mi}(T)} y^{\text{fd}(T)} z^{\text{snd}(T)} \bar{x}^{\text{mp}(T)} \bar{y}^{\text{fnd}(T)} \bar{z}^{\text{sd}(T)} = \langle W | N_{i_1} \dots N_{i_n} | V \rangle,$$

where the sum is over $T \in \mathcal{S}_n$ of profile (i_1, \dots, i_n) .

Proof. We follow the same scheme as in Proposition 4.5, and use Proposition 5.3. Here, the surjective pretableaux of a given profile are in bijection with surjective tableaux of a particular shape. Rather than giving a formal detailed proof, we sketch how to understand the matrices N_1 through N_6 , having in mind the proof of Proposition 5.3 and the way surjective tableaux are build recursively.

- The k th co-corner and k th corner correspond to the k th factor BBA in the word $(BBA)^n$. If these are empty (*i.e.* $i_k = 1$), we can remove the two cells and replace the factor BBA with ABB . With the terms xI and $\bar{x}I$ as seen before, we see that this case correspond to the matrix $N_1 = A(B + xI)(B + \bar{x}I)$.
- If $i_k = 2$, *i.e.* the k th co-corner and k th corner both contain a \times , we can remove the two columns containing these \times , and remove the factor BB in BBA . But we need to distinguish two cases, depending on wether there is a third \times in the same row or not, and if there is not, we also remove the row. This gives a factor $A + I$, and there is a weight $y\bar{z}$ because of the doubled \times in the corner and co-corner. Hence this case correspond to the matrix $N_2 = y\bar{z}(A + I)$.
- If $i_k = 3$, *i.e.* the k th co-corner contains a non-doubled \times , we can remove its row and column, so the k th factor BBA becomes a $B + \bar{x}I$. There is a weight z for the non-doubled \times in the co-corner. Hence this case gives the matrix $N_3 = z(B + \bar{x}I)$.
- If $i_k = 4$, the difference with the previous case is that the \times in the k th co-corner is doubled, so we do not remove its row. So there remains a factor A , and there is a weight \bar{z} instead of z . Hence this case gives the matrix $N_4 = \bar{z}A(B + \bar{x}I)$.
- If $i_k = 5$, this is similar to the case when $i_k = 3$. But the weight is \bar{y} instead of z for the non-doubled \times in the corner, and there remains a column of odd index so this gives a factor $B + xI$ instead of $B + \bar{x}I$. Hence this case gives the matrix $N_5 = \bar{y}(B + xI)$.
- If $i_k = 6$, this is similar to the case when $i_k = 4$. But the weight is y instead of \bar{z} for the doubled \times in the corner, and there remains a column of odd index so this gives a factor $B + xI$ instead of $B + \bar{x}I$. Hence this case gives the matrix $N_6 = yA(B + xI)$.

When we form the product $N_{i_1} \dots N_{i_n}$, it is clear that the matrix N_{i_k} will impose the conditions on the k th co-corner and corner, and hence $\langle W | N_{i_1} \dots N_{i_n} | V \rangle$ is the generating function for elements in \mathcal{S}_n of profile (i_1, \dots, i_n) . \square

Now, we can prove Proposition 5.4.

Proof. We have $N = \sum_{i=1}^6 N_i$, hence using Lemma 5.6:

$$\begin{aligned} \langle W|N^n|V \rangle &= \sum_{(i_1, \dots, i_n) \in \{1, \dots, 6\}^n} \langle W|N_{i_1} \dots N_{i_n}|V \rangle \\ &= \sum_{T \in \mathcal{S}_n} x^{\text{mi}(T)} y^{\text{fd}(T)} z^{\text{snd}(T)} \bar{x}^{\text{mp}(T)} \bar{y}^{\text{fnd}(T)} \bar{z}^{\text{sd}(T)} = \Gamma_{n+1}. \end{aligned}$$

This completes the proof. \square

From the definitions of B and A in (26), the matrix N defined in (27) can be calculated explicitly and we obtain the following statement.

Proposition 5.7. *The matrix $N = (N_{i,j})_{i,j \in \mathbb{N}}$ is tridiagonal, and such that for any $i \geq 0$ we have $N_{i,i} = b_i$ and $N_{i,i+1}N_{i+1,i} = \lambda_{i+1}$.*

Proof. Straightforward calculations show that $N_{i,j} = 0$ if $|i - j| > 1$, $N_{i,i} = b_i$, and:

$$(29) \quad N_{i,i+1} = (i+1)(z + \bar{y} + i), \quad N_{i+1,i} = (x + \bar{z} + i)(y + \bar{x} + i).$$

This gives indeed $N_{i,i+1}N_{i+1,i} = \lambda_{i+1}$. \square

As in the case of alternative tableaux in the previous section, the previous two propositions means that the continued fraction expansion given in (4) can be derived from the combinatorial interpretation in terms of escaliers, and the matrix Ansatz for escaliers given in Proposition 5.3.

Remark 5.8. Observe that $N \neq M$ since the non-diagonal coefficients are not the same, but the two matrices are equal after a permutation of the variables $(x, y, z, \bar{x}, \bar{y}, \bar{z})$. However, there is *a priori* no simple way to link the matrices M_i 's with the N_i 's, so we tend to think that despite the similarities in the method, there are two really different ways to obtain the continued fraction from a combinatorial model using the matrix Ansatz approach.

It is natural to ask if there is a bijection between \mathcal{T}_n and \mathcal{S}_n preserving the six statistics for Γ_n . From the fact that the recurrence relation is checked on both sets, in theory it might be possible to describe recursively such a bijection. It would be quite interesting to give a better answer to this question by providing a direct bijection between our alternative tableaux and Dumont's escaliers.

CONCLUSION

Our main result is the new combinatorial interpretation of Γ_n with the alternative tableaux. We obtain two new proofs of Dumont's conjecture, with the same method applied to the two different combinatorial interpretations of Γ_n . What it is interesting about these proofs is that they fit in the general framework settled in [1], linking J-fractions, operators satisfying certain commutation relations, and combinatorial objects.

ACKNOWLEDGEMENT

I thank Dominique Dumont, Dominique Foata, Jiang Zeng, Xavier Viennot and Sylvie Corteel for their conversation and encouragements.

REFERENCES

- [1] S. Corteel, M. Josuat-Vergès and L.K. Williams, Matrix Ansatz, orthogonal polynomials and permutations, to appear in Adv. Appl. Math. (2010).
- [2] S. Corteel and L.K. Williams, Tableaux combinatorics for the asymmetric exclusion process, Adv. in Appl. Math. 39(3) (2007), 293–310.
- [3] B. Derrida, M.R. Evans, V. Hakim and V. Pasquier, Exact solution of a 1D asymmetric exclusion model using a matrix formulation, J. Phys. A 26 (1993), 1493–1517.
- [4] D. Dumont, Conjectures sur des symétries ternaires liées au nombres de Genocchi, Discrete Math. 139 (1995), 469–472.
- [5] D. Dumont and D. Foata, Une propriété de symétrie des nombres de Genocchi, Bull. Soc. Math. France 104 (1976), 433–451.
- [6] P. Flajolet, Combinatorial aspects of continued fractions, Discrete Math. 41 (1982), 145–153.
- [7] I. Gessel and J. Zeng, unpublished.
- [8] R. Koekoek and R.F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue, Delft University of Technology, Report no. 98–17 (1998).
- [9] P. Nadeau, The structure of alternative tableaux, arXiv:0908.4050v2 [math.CO].
- [10] A. Randrianarivony, Polynômes de Dumont-Foata généralisés, Sémin. Lothar. Combin. 32 (1994), Article B32d, 12 pp.
- [11] A. Randrianarivony and J. Zeng, Sur une extension des nombres d’Euler et les records des permutations alternantes, J. Combin. Theory Ser. A 68 (1994), 86–99.
- [12] X.G. Viennot, Interprétations combinatoires des nombres d’Euler et de Genocchi, séminaire de théorie des nombres de l’Université Bordeaux I, 1981.
- [13] X.G. Viennot, Alternative tableaux, permutations and partially asymmetric exclusion process, talk in Isaac Newton institute, April 2008.
<http://www.newton.ac.uk/webseminars/pg+ws/2008/csm/csmw04/0423/viennot/>
- [14] J. Ziang, Sur quelques propriétés de symétrie des nombres de Genocchi, Discrete Math. 153 (1996), 319–333.

LRI, UNIVERSITÉ PARIS-SUD, BÂT. 490, 91405 ORSAY, FRANCE
E-mail address: josuat@lri.fr