


A phase transition in block-weighted random maps

Journées Combinatoires de Bordeaux
5 février 2024


Zéphyr Salvy (he/they)

Planar maps

Planar map \mathfrak{m} = embedding on the sphere of a connected planar graph, considered up to homeomorphisms




Planar map = planar graph +
cyclic order on neighbours



- **Rooted** planar map = map endowed with a marked oriented edge (represented by an arrow);
- **Size** $|\mathfrak{m}|$ = number of edges;
- **Corner** (does not exist for graphs !) = space between an oriented edge and the next one for the trigonometric order.

Universality results for planar maps

- Enumeration: $\kappa \rho^{-n} n^{-5/2}$ [Tutte 1963];
- Distance between vertices: $n^{1/4}$ [Chassaing, Schaeffer 2004];
- Scaling limit: Brownian sphere for quadrangulations [Le Gall 2013, Miermont 2013] and general maps [Bettinelli, Jacob, Miermont 2014];



Brownian Sphere \mathcal{S}_e




Universality results for planar maps

- Enumeration: $\kappa \rho^{-n} n^{-5/2}$ [Tutte 1963];
- Distance between vertices: $n^{1/4}$ [Chassaing, Schaeffer 2004];
- Scaling limit: Brownian sphere for quadrangulations [Le Gall 2013, Miermont 2013] and general maps [Bettinelli, Jacob, Miermont 2014];
- Universality:
 - Same enumeration [Drmotá, Noy, Yu 2020];
 - Same scaling limit, e.g. for triangulations & $2q$ -angulations [Le Gall 2013], simple quadrangulations [Addario-Berry, Albenque 2017].

Universality results for plane trees

- Enumeration: $\kappa \rho^{-n} n^{-3/2}$;
- Distance between vertices: $n^{1/2}$ [Flajolet, Odlyzko 1982];
- Scaling limit: Brownian tree [Aldous 1993, Le Gall 2006];



Universality results for plane trees

- Enumeration: $\kappa \rho^{-n} n^{-3/2}$;
- Distance between vertices: $n^{1/2}$ [Flajolet, Odlyzko 1982];
- Scaling limit: Brownian tree [Aldous 1993, Le Gall 2006];
- Universality:
 - Same enumeration,
 - Same scaling limit, even for some classes of **maps**; e.g. outerplanar maps [Caraceni 2016], maps with a boundary of size $\gg n^{1/2}$ [Bettinelli 2015].


Models with (very) constrained boundaries

Motivation



Inspired by [Bonzom 2016].

Two rich situations with universality results:

Planar maps




Plane trees



Brownian Sphere \mathcal{S}_e

Interpolating model?



Brownian Tree \mathcal{T}_e

Model definition


2-connected = two vertices must be removed to disconnect.

Block = maximal (for inclusion) 2-connected submap.

Model definition

2-connected = two vertices must be removed to disconnect.


Block = maximal (for inclusion) 2-connected submap.



Model definition

2-connected = two vertices must be removed to disconnect.

Block = maximal (for inclusion) 2-connected submap.




Condensation phenomenon: a large block concentrates a macroscopic part of the mass
[Banderier, Flajolet, Schaeffer, Soria 2001; Jonsson, Stefánsson 2011].


Model definition

2-connected = two vertices must be removed to disconnect.

Block = maximal (for inclusion) 2-connected submap.



Condensation phenomenon: a large block concentrates a macroscopic part of the mass [Banderier, Flajolet, Schaeffer, Soria 2001; Jonsson, Stefánsson 2011].




Only small blocks.


Model definition

2-connected = two vertices must be removed to disconnect.

Block = maximal (for inclusion) 2-connected submap.



Condensation phenomenon: a large block concentrates a macroscopic part of the mass [Banderier, Flajolet, Schaeffer, Soria 2001; Jonsson, Stefánsson 2011].



Only small blocks.

Interpolating model using blocks!

Outline of the talk

A phase transition in block-weighted random maps

I. Model

II. Block tree of a map and its applications

Interlude. Quadrangulations

III. Scaling limits

IV. Extension to other families of maps


V. Extension to tree-rooted maps

VI. Perspectives

with William
Fleurat



with Marie
Albenque & Éric
Fusy



I. Model

Model

Inspired by [Bonzom 2016].

Goal: parameter that affects the typical number of blocks.


We choose: $\mathbb{P}_{n,u}(\mathfrak{m}) = \frac{u^{\#\text{blocks}(\mathfrak{m})}}{Z_{n,u}}$ where

$u > 0,$
 $\mathcal{M}_n = \{\text{maps of size } n\},$
 $\mathfrak{m} \in \mathcal{M}_n,$
 $Z_{n,u} = \text{normalisation.}$

- $u = 1$: uniform distribution on maps of size n ;
- $u \rightarrow 0$: minimising the number of blocks (=2-connected maps);
- $u \rightarrow \infty$: maximising the number of blocks (= trees!).


Given u , asymptotic behaviour when $n \rightarrow \infty$?

$u = 1$



$n \approx 55\,000$

$u = 8/5$




$n \approx 55\,000$

$u = 9/5$



$n \approx 80\,000$

$$u = 5/2$$



$$n \approx 75\,000$$

$u = 5$



$n \approx 50\,000$

Phase transition

Theorem [Fleurat, S. 23] Model exhibits a phase transition at $u = 9/5$. When $n \rightarrow \infty$:

- Subcritical phase $u < 9/5$: “general map phase” one huge block;
- Critical phase $u = 9/5$: a few large blocks;
- Supercritical phase $u > 9/5$: “tree phase” only small blocks.

We obtain explicit results on enumeration, size of blocks and scaling limits in each case.

→ *A phase transition in block-weighted random maps*
W. Fleurat & Z. S., *Electronic Journal of Probability*, 2024


Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
Enumeration			
Size of - the largest block - the second one			
Scaling limit of M_n			

II. Block tree of a map and its applications



Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]





Decomposition of a map into blocks

Inspiration from [Tutte 1963]




Decomposition of a map into blocks

Inspiration from [Tutte 1963]





Decomposition of a map into blocks

Inspiration from [Tutte 1963]





Decomposition of a map into blocks

Inspiration from [Tutte 1963]





Decomposition of a map into blocks

Inspiration from [Tutte 1963]





Decomposition of a map into blocks

Inspiration from [Tutte 1963]



Decomposition of a map into blocks

Inspiration from [Tutte 1963]



$$M(z) = B(zM^2(z))$$



GS of 2-connected maps



Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#blocks(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



$$M(z) = B(zM^2(z))$$

GS of 2-connected maps


With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
Enumeration <small>[Bonzom 2016]</small>	$\rho(u)^{-n} n^{-5/2}$	$\rho(u)^{-n} n^{-5/3}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one			
Scaling limit of M_n			

Decomposition of a map into blocks



Inspiration from [Tutte 1963]



Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-

Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-

Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-



Berry 2019].
 GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-

Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-

Berry 2019].



GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



⇒ Underlying block tree structure, made explicit by [Addario-


Berry 2019].
 GS of 2-connected maps

With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks

$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]





With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Decomposition of a map into blocks


$$M(z, u) = \sum_{\mathfrak{m} \in \mathcal{M}} z^{|\mathfrak{m}|} u^{\#\text{blocks}(\mathfrak{m})}$$

Inspiration from [Tutte 1963]



With a weight u on blocks: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Properties of the block tree



- \mathfrak{m} is entirely determined by $T_{\mathfrak{m}}$ and $(\mathfrak{b}_v, v \in T_{\mathfrak{m}})$ where \mathfrak{b}_v is the block of \mathfrak{m} represented by v in $T_{\mathfrak{m}}$;
- Internal node (with $2k$ children) of $T_{\mathfrak{m}} \leftrightarrow$ block of \mathfrak{m} of size k .

T_{M_n} gives the block sizes of a random map M_n .

Galton-Watson trees for map blocks

μ -Galton-Watson tree : random tree where the number of children of each node is given by μ independently, with μ = probability law on \mathbb{N} .

Galton-Watson trees for map blocks

μ -Galton-Watson tree : random tree where the number of children of each node is given by μ independently, with μ = probability law on \mathbb{N} .

Theorem [Fleurat, S. 23]

$u > 0$

If $M_n \hookrightarrow \mathbb{P}_{n,u}$ then there exists an (explicit) $y = y(u)$ s.t.

T_{M_n} has the law of a Galton-Watson tree of reproduction

law $\mu^{y,u}$ conditioned to be of size $2n$, with


$$\mu^{y,u}(\{2k\}) = \frac{B_k y^k u^{1_{k \neq 0}}}{uB(y) + 1 - u}.$$

Phase transition


When is $\mu^{y,u}$ critical? ($= \mathbb{E}(\mu) = 1$?)

$$\mathbb{E}(\mu^{y,u}) = 1 \Leftrightarrow u = \frac{1}{2yB'(y) - B(y) + 1}$$

covers $[9/5, +\infty)$ when y covers $(0, \rho_B = 4/27]$.



Phase transition



Largest blocks?

- Degrees of T_{M_n} give the block sizes of the map M_n ;
- Largest degrees of a Galton-Watson tree are well-known [Janson 2012].

Rough intuition


	$u < 9/5$	$u = 9/5$	$u > 9/5$
$\mu^{y(u),u}(\{2k\})$	$\sim c_u k^{-5/2}$		$\sim c_u \pi_u^k k^{-5/2}$
Galton-Watson tree	subcritical	critical	

Dichotomy between situations:

- Subcritical: condensation, cf [Jonsson Stefánsson 2011];
- Supercritical: behaves as maximum of independent variables.

Size $L_{n,k}$ of the k -th largest block

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
$L_{n,1}$	$\sim (1 - \mathbb{E}(\mu^{4/27,u}))n$ [Stufler 2020]		
$L_{n,2}$	$\Theta(n^{2/3})$ [Stufler 2020]	$\Theta(n^{2/3})$	$\frac{\ln(n)}{2 \ln\left(\frac{4}{27y}\right)} - \frac{5 \ln(\ln(n))}{4 \ln\left(\frac{4}{27y}\right)} + O(1)$





Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
Enumeration [Bonzom 2016]	$\rho(u)^{-n} n^{-5/2}$	$\rho(u)^{-n} n^{-5/3}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one	$\sim (1 - \mathbb{E}(\mu^{4/27,u}))n$ $\Theta(n^{2/3})$ [Stufler 2020]	$\Theta(n^{2/3})$	$\frac{\ln(n)}{2 \ln\left(\frac{4}{27y}\right)} - \frac{5 \ln(\ln(n))}{4 \ln\left(\frac{4}{27y}\right)} + O(1)$
Scaling limit of M_n			

Interlude: quadrangulations

Quadrangulations

Def: map with all faces of degree 4.




Simple quadrangulation = no multiple edges.


Size $|\mathfrak{q}|$ = number of *faces*.

$$|V(\mathfrak{q})| = |\mathfrak{q}| + 2, \quad |E(\mathfrak{q})| = 2|\mathfrak{q}|.$$


Construction of a quadrangulation from a simple core





Construction of a quadrangulation from a simple core



Construction of a quadrangulation from a simple core



Block tree for a quadrangulation





With a weight u on blocks: $Q(z, u) = uS(zQ^2(z, u)) + 1 - u$



Remember: $M(z, u) = uB(zM^2(z, u)) + 1 - u$

Tutte's bijection

Map




Quadrangulation



[Tutte 1963]

Tutte's bijection for 2-connected maps







Cut vertex \Rightarrow multiple edge

2-connected maps \Leftrightarrow simple quadrangulations

[Brown 1965]

Block trees under Tutte's bijection



Implications on results

We choose: $\mathbb{P}_{n,u}(\mathbf{q}) = \frac{u^{\#blocks(\mathbf{q})}}{Z_{n,u}}$ where

$u > 0,$
 $\mathcal{Q}_n = \{\text{quadrangulations of size } n\},$
 $\mathbf{q} \in \mathcal{Q}_n,$
 $Z_{n,u} = \text{normalisation.}$

Results on the size of (2-connected) blocks can be transferred immediately for quadrangulations and their simple blocks.


Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
Enumeration <small>[Bonzom 2016] for 2-c case</small>	$\rho(u)^{-n} n^{-5/2}$	$\rho(u)^{-n} n^{-5/3}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one	$\sim (1 - \mathbb{E}(\mu^{4/27,u}))n$ $\Theta(n^{2/3})$ <small>[Stufler 2020]</small>	$\Theta(n^{2/3})$	$\frac{\ln(n)}{2 \ln\left(\frac{4}{27y}\right)} - \frac{5 \ln(\ln(n))}{4 \ln\left(\frac{4}{27y}\right)} + O(1)$
Scaling limit of M_n			

III. Scaling limits

Scaling limits

Convergence of the whole object considered as a metric space (with the graph distance), after renormalisation.



$M_n \hookrightarrow \mathbb{P}_{n,u}$ (map or quadrangulation)

What is the limit of the sequence of metric spaces $((M_n, d/n^?)_{n \in \mathbb{N}}$?

(Convergence for Gromov-Hausdorff topology)

Scaling limit of supercritical and critical maps


Lemma For $M_n \hookrightarrow \mathbb{P}_{n,u'}$

- If $u > 9/5$,

$$\frac{c_3(u)}{n^{1/2}} T_{M_n} \rightarrow \mathcal{T}_e.$$

- If $u = 9/5$,

$$\frac{c_2}{n^{1/3}} T_{M_n} \rightarrow \mathcal{T}_{3/2}.$$



Brownian Tree \mathcal{T}_e

Stable Tree $\mathcal{T}_{3/2}$

Scaling limit of supercritical and critical maps

Lemma For $M_n \hookrightarrow \mathbb{P}_{n,u}$

- If $u > 9/5$,

$$\frac{c_3(u)}{n^{1/2}} T_{M_n} \rightarrow \mathcal{T}_e.$$

- If $u = 9/5$,

$$\frac{c_2}{n^{1/3}} T_{M_n} \rightarrow \mathcal{T}_{3/2}.$$

Proof Known scaling limits of critical Galton-Watson trees

- with finite variance [Aldous 1993, Le Gall 2006];
- infinite variance and polynomial tails [Duquesne 2003].

Scaling limit of supercritical and critical maps

Theorem For $M_n \hookrightarrow \mathbb{P}_{n,u'}$

- [Stufler 2020] If $u > 9/5$,


$$\frac{c_3(u)}{n^{1/2}} T_{M_n} \rightarrow \mathcal{T}_e$$

$$\frac{C_3(u)}{n^{1/2}} M_n \rightarrow \mathcal{T}_e.$$


- [Fleurat, S. 23] If $u = 9/5$,

$$\frac{c_2}{n^{1/3}} T_{M_n} \rightarrow \mathcal{T}_{3/2}$$

$$\frac{C_2}{n^{1/3}} M_n \rightarrow \mathcal{T}_{3/2}.$$



Brownian Tree \mathcal{T}_e



Stable Tree $\mathcal{T}_{3/2}$

Scaling limit of supercritical and critical maps

Theorem For $M_n \hookrightarrow \mathbb{P}_{n,u}$

- [Stufler 2020] If $u > 9/5$,

$$\frac{c_3(u)}{n^{1/2}} T_{M_n} \rightarrow \mathcal{T}_e$$

$$\frac{C_3(u)}{n^{1/2}} M_n \rightarrow \mathcal{T}_e.$$

- [Fleurat, S. 23] If $u = 9/5$,


$$\frac{c_2}{n^{1/3}} T_{M_n} \rightarrow \mathcal{T}_{3/2}$$

$$\frac{C_2}{n^{1/3}} M_n \rightarrow \mathcal{T}_{3/2}.$$

Proof Distances in M_n behave like distances in T_{M_n} !

Supercritical and critical cases

Difficult part = show that distances in M_n behave like distances in T_{M_n} .



Let $\kappa = \mathbb{E}$ ("diameter" bipointed block). By a "law of large numbers"-type argument

$$d_{M_n}(e_1, e_2) \simeq \kappa d_{T_{M_n}}(e_1, e_2).$$


Difficult for the critical case => large deviation estimates

$u = 9/5$



$n \approx 80\,000$

$$u = 5/2$$



$$n \approx 75\,000$$

$u = 5$



$n \approx 50\,000$

Scaling limits of subcritical maps


Theorem [Fleurat, S. 23] If $u < 9/5$, for $M_n \hookrightarrow \mathbb{P}_{n,u}$ and denoting $B(M_n)$ its largest block:

$$d_{GHP} \left(\frac{C_1(u)}{n^{1/4}} M_n, \frac{1}{n^{1/4}} B(M_n) \right) \rightarrow 0.$$

Brownian Sphere \mathcal{S}_e

So, if $cn^{-1/4} B_n \rightarrow \mathcal{S}_e$, then

$$\frac{C_1(u)}{cn^{1/4}} M_n \rightarrow \mathcal{S}_e.$$



See [Addario-Berry, Wen 2019] for a similar result and method.

Scaling limits of subcritical maps


Theorem [Fleurbaey, S. 23] If $u < 9/5$, for $Q_n \hookrightarrow \mathbb{P}_{n,u}$ a quadrangulation:

$$\frac{C_1(u)}{n^{1/4}} Q_n \rightarrow \mathcal{S}_e.$$


Moreover, Q_n and its simple core converge jointly to the same Brownian sphere.

Proof

- Previous theorem;
- Scaling limit of uniform simple quad. rescaled by $n^{1/4}$ = Brownian sphere [Addario-Berry Albenque 2017].



Subcritical case



Diameter of a decoration \leq blocks to cross \times max diameter of blocks


$$\leq \text{diam}(T_{M_n}) \times (O(n^{2/3}))^{1/4+\delta} = \text{diam}(T_{M_n}) \times O(n^{1/6+\delta})$$

T_{M_n} is a subcritical Galton-Watson tree

$$= O(n^{1/6+2\delta}) = o(n^{1/4}).$$

[Chapuy Fusy Giménez Noy 2015]

Subcritical case



Decorations = groups of smaller blocks

Diameters of decorations = $o(n^{1/4})$.

Diameter of a decoration \leq blocks to cross \times max diameter of blocks


$$\leq \text{diam}(T_{M_n}) \times (O(n^{2/3}))^{1/4+\delta} = \text{diam}(T_{M_n}) \times O(n^{1/6+\delta})$$

T_{M_n} is a subcritical Galton-Watson tree

$$= O(n^{1/6+2\delta}) = o(n^{1/4}).$$


[Chapuy Fusy Giménez Noy 2015]

Subcritical case




The scaling limit of M_n (rescaled by $n^{1/4}$) is the scaling limit of uniform blocks!

$u = 1$






$n \approx 55\,000$

$u = 8/5$



$n \approx 55\,000$

Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < 9/5$	$u = 9/5$	$u > 9/5$
Enumeration [Bonzom 2016]	$\rho(u)^{-n} n^{-5/2}$	$\rho(u)^{-n} n^{-5/3}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one	$\sim (1 - \mathbb{E}(\mu^{4/27,u}))n$ $\Theta(n^{2/3})$ [Stufler 2020]	$\Theta(n^{2/3})$	$\frac{\ln(n)}{2 \ln\left(\frac{4}{27y}\right)} - \frac{5 \ln(\ln(n))}{4 \ln\left(\frac{4}{27y}\right)} + O(1)$
Scaling limit of M_n	$\frac{C_1(u)}{n^{1/4}} M_n \rightarrow \mathcal{S}_e$ 	$\frac{C_2}{n^{1/3}} M_n \rightarrow \mathcal{T}_{3/2}$ 	$\frac{C_3(u)}{n^{1/2}} M_n \rightarrow \mathcal{T}_e$ [Stufler 2020] 

Assuming the convergence of 2-connected maps towards the Brownian sphere

IV. Extension to other families of maps

Extension to other models

[Banderier, Flajolet, Schaeffer, Soria 2001]:

TABLE 3. Composition schemas, of the form $\mathcal{M} = \mathcal{C} \circ \mathcal{H} + \mathcal{D}$, except the last one where $\mathcal{M} = (1 + \mathcal{M}) \times (\mathcal{C} \circ \mathcal{H})$.

maps, $M(z)$	cores, $C(z)$	submaps, $H(z)$	coreless, $D(z)$
all, $M_1(z)$	bridgeless, $M_2(z)$ or loopless	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$
loopless $M_2(z)$	simple $M_3(z)$	$z(1 + M)$	–
all, $M_1(z)$	nonsep., $M_4(z)$	$z(1 + M)^2$	–
nonsep. $M_4(z) - z$	nonsep. simple $M_5(z)$	$z(1 + M)$	–
nonsep. $M_4(z)/z - 2$	3-connected $M_6(z)$	M	$z + 2M^2/(1 + M)$
bipartite, $B_1(z)$	bip. simple, $B_2(z)$	$z(1 + M)$	–
bipartite, $B_1(z)$	bip. bridgeless, $B_3(z)$	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$
bipartite, $B_1(z)$	bip. nonsep., $B_4(z)$	$z(1 + M)^2$	–
bip. nonsep., $B_4(z)$	bip. ns. simpl, $B_5(z)$	$z(1 + M)$	–
singular tri., $T_1(z)$	triang., $z + zT_2(z)$	$z(1 + M)^3$	–
triangulations, $T_2(z)$	irreducible tri., $T_3(z)$	$z(1 + M)^2$	–

Extension to other models

[Banderier, Flajolet, Schaeffer, Soria 2001]:

TABLE 3. Composition schemas, of the form $\mathcal{M} = \mathcal{C} \circ \mathcal{H} + \mathcal{D}$, except the last one where $\mathcal{M} = (1 + \mathcal{M}) \times (\mathcal{C} \circ \mathcal{H})$.

maps, $M(z)$	cores, $C(z)$	submaps, $H(z)$	coreless, $D(z)$	u_C
all, $M_1(z)$	bridgeless, $M_2(z)$ or loopless	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$	
loopless $M_2(z)$	simple $M_3(z)$	$z(1 + M)$	–	81/17
all, $M_1(z)$	nonsep., $M_4(z)$	$z(1 + M)^2$	–	9/5
nonsep. $M_4(z) - z$	nonsep. simple $M_5(z)$	$z(1 + M)$	–	135/7
nonsep. $M_4(z)/z - 2$	3-connected $M_6(z)$	M	$z + 2M^2/(1 + M)$	
bipartite, $B_1(z)$	bip. simple, $B_2(z)$	$z(1 + M)$	–	36/11
bipartite, $B_1(z)$	bip. bridgeless, $B_3(z)$	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$	
bipartite, $B_1(z)$	bip. nonsep., $B_4(z)$	$z(1 + M)^2$	–	52/27
bip. nonsep., $B_4(z)$	bip. ns. simpl, $B_5(z)$	$z(1 + M)$	–	68/3
singular tri., $T_1(z)$	triang., $z + zT_2(z)$	$z(1 + M)^3$	–	16/7
triangulations, $T_2(z)$	irreducible tri., $T_3(z)$	$z(1 + M)^2$	–	64/37

→ *Unified study of the phase transition for block-weighted random planar maps* Z. Salvy (Eurocomb'23)

Statement of the results

Theorem [S. 23] Model of the preceding table without coreless maps exhibits a **phase transition** at some **explicit** u_C .

When $n \rightarrow \infty$:

- Subcritical phase $u < u_C$: “**general map phase**” one huge block;
- Critical phase $u = u_C$: a few large blocks;
- Supercritical phase $u > u_C$: “**tree phase**” only small blocks.


We obtain **explicit** results on enumeration and size of blocks in each case.

V. Extension to tree-rooted maps

Decorated maps are interesting

From a theoretical physics point of view:


- Before: “pure gravity” case (nothing happens on the surface);
- Now: decorated map (things happen! new behaviours! excitement!).



Tree-rooted maps



= (rooted planar) maps endowed with a spanning tree.




$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n$$

[Mullin 67]

We want to study block-weighted tree-rooted maps.

Block decomposition of tree-rooted maps

The decomposition of maps into blocks extends into a decomposition of tree-rooted maps into tree-rooted blocks.



$$M(z) = B(zM^2(z))$$

GS of 2-connected tree-rooted maps

Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$

Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$

- $[z^n]M(z) \sim \frac{4}{\pi} \times 16^n \times n^{-3};$

Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$


$$\bullet [z^n]M(z) \sim \frac{4}{\pi} \times 16^n \times n^{-3}; \quad \bullet \rho_M = \frac{1}{16};$$

Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$

$$\bullet [z^n]M(z) \sim \frac{4}{\pi} \times 16^n \times n^{-3}; \quad \bullet \rho_M = \frac{1}{16};$$

$$\bullet M(\rho_M) = 8 - \frac{64}{3\pi} \simeq 1.2 \text{ so } M \text{ is not algebraic...}$$


$$P(z, M(z)) = 0$$


Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$

$$\bullet [z^n]M(z) \sim \frac{4}{\pi} \times 16^n \times n^{-3}; \quad \bullet \rho_M = \frac{1}{16};$$

$$\bullet M(\rho_M) = 8 - \frac{64}{3\pi} \simeq 1.2 \text{ so } M \text{ is not algebraic...}$$



$$P(z, M(z)) = 0$$

An arrow points from the text above to this equation, which is crossed out with a large pink 'X', indicating that this equation does not hold for the series M(z).


Tree-rooted maps are not so nice

$$M(z) = \sum_{n \geq 0} \text{Cat}_n \text{Cat}_{n+1} z^n \text{ so}$$


$$\bullet [z^n]M(z) \sim \frac{4}{\pi} \times 16^n \times n^{-3}; \quad \bullet \rho_M = \frac{1}{16};$$

$$\bullet M(\rho_M) = 8 - \frac{64}{3\pi} \simeq 1.2 \text{ so } M \text{ is not algebraic...}$$

• Fortunately, it is still *D-finite*

$$P(z, M(z)) = 0$$


$$P_0(z) \frac{\partial^2 M}{\partial z^2}(z) + P_1(z) \frac{\partial M}{\partial z}(z) + P_2(z) M(z) + P_3(z) = 0.$$



2-connected tree-rooted maps are naughty


Using $M(z) = B(zM^2(z))$ and the properties of M , we show

2-connected tree-rooted maps are naughty

Using $M(z) = B(zM^2(z))$ and the properties of M , we show

- $\rho_B = \rho_M M^2(\rho_M) = \frac{4(3\pi - 8)^2}{9\pi^2} \approx 0.091$

is not algebraic so B is not D -finite



2-connected tree-rooted maps are naughty


Using $M(z) = B(zM^2(z))$ and the properties of M , we show

- $\rho_B = \rho_M M^2(\rho_M) = \frac{4(3\pi - 8)^2}{9\pi^2} \approx 0.091$

is not algebraic so B is not D -finite

- B is D -algebraic

$$P\left(\frac{\partial^2 B}{\partial y^2}(y), \frac{\partial B}{\partial y}(y), B(y), y\right) = 0.$$



Enumeration of 2-connected tree-rooted maps


Using $M(z) = B(zM^2(z))$ and the properties of M , we show

Theorem [Albenque, Fusy, S. 24+]

$$[y^n]B(y) \sim \frac{4(3\pi - 8)^3}{27\pi(4 - \pi)^3} \times \rho_B^{-n} \times n^{-3}.$$

Block decomposition of tree-rooted maps

The decomposition of maps into blocks extends into a decomposition of tree-rooted maps into tree-rooted blocks.




$$M(z) = B(zM^2(z))$$

GS of 2-connected tree-rooted maps

Block decomposition of tree-rooted maps

The decomposition of maps into blocks extends into a decomposition of tree-rooted maps into tree-rooted blocks.



$$M(z, u) = uB(zM^2(z, u)) + 1 - u$$

Phase transition

Theorem [Albenque, Fusy, S. 24+] Model exhibits a phase

transition at $u_C = \frac{9\pi(4 - \pi)}{420\pi - 81\pi^2 - 512} \simeq 3.02$.

When $n \rightarrow \infty$:

- Subcritical phase $u < u_C$: “general tree-rooted map phase” one huge block;
- Critical phase $u = u_C$: a few large blocks;
- Supercritical phase $u > u_C$: “tree phase” only small blocks.

We obtain results on enumeration, size of blocks and scaling limits in each case.

Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < u_C$	$u = u_C$	$u > u_C$
Enumeration			
Size of - the largest block - the second one			
Scaling limit of M_n			

Results


For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < u_C$	$u = u_C$	$u > u_C$
Enumeration	$\rho(u)^{-n} n^{-3}$	$\rho(u)^{-n} n^{-3/2} \ln(n)^{-1/2}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one			
Scaling limit of M_n			

Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < u_C$	$u = u_C$	$u > u_C$
Enumeration	$\rho(u)^{-n} n^{-3}$	$\rho(u)^{-n} n^{-3/2} \ln(n)^{-1/2}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one	$\sim (1 - \mathbb{E}(\mu^{\rho_B, u}))n$ $\Theta(n^{1/2})$	$\Theta(n^{1/2})$	$\frac{\ln(n)}{\ln\left(\frac{\rho_B}{y(u)}\right)} - \frac{3 \ln(\ln(n))}{\ln\left(\frac{\rho_B}{y(u)}\right)} + O(1)$
Scaling limit of M_n			

Results

For $M_n \hookrightarrow \mathbb{P}_{n,u}$	$u < u_C$	$u = u_C$	$u > u_C$
Enumeration	$\rho(u)^{-n} n^{-3}$	$\rho(u)^{-n} n^{-3/2} \ln(n)^{-1/2}$	$\rho(u)^{-n} n^{-3/2}$
Size of - the largest block - the second one	$\sim (1 - \mathbb{E}(\mu^{\rho_{B,u}}))n$ $\Theta(n^{1/2})$	$\Theta(n^{1/2})$	$\frac{\ln(n)}{\ln\left(\frac{\rho_B}{y(u)}\right)} - \frac{3 \ln(\ln(n))}{\ln\left(\frac{\rho_B}{y(u)}\right)} + O(1)$
Scaling limit of M_n	?	$\frac{C_2}{n^{1/2} \ln(n)^{1/2}} M_n \rightarrow \mathcal{T}_e$	$\frac{C_3(u)}{n^{1/2}} M_n \rightarrow \mathcal{T}_e$



Interlude: tree-rooted quadrangulations

CANCELLED

**Interlude: tree-rooted
quadragulations**

$M(z) = Q(z)$ does not hold!

VI. Perspectives

Extension to more involved decompositions

- For maps : maps into loopless blocks, 2-connected maps into 3-connected blocks;

Extension to more involved decompositions

- For maps : maps into loopless blocks, 2-connected maps into 3-connected blocks;

TABLE 3. Composition schemas, of the form $\mathcal{M} = \mathcal{C} \circ \mathcal{H} + \mathcal{D}$, except the last one where $\mathcal{M} = (1 + \mathcal{M}) \times (\mathcal{C} \circ \mathcal{H})$.

maps, $M(z)$	cores, $C(z)$	submaps, $H(z)$	coreless, $D(z)$
all, $M_1(z)$	bridgeless, or loopless $M_2(z)$	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$
loopless $M_2(z)$	simple $M_3(z)$	$z(1 + M)$	–
all, $M_1(z)$	nonsep., $M_4(z)$	$z(1 + M)^2$	–
nonsep. $M_4(z) - z$	nonsep. simple $M_5(z)$	$z(1 + M)$	–
nonsep. $M_4(z)/z - 2$	3-connected $M_6(z)$	M	$z + 2M^2/(1 + M)$
bipartite, $B_1(z)$	bip. simple, $B_2(z)$	$z(1 + M)$	–
bipartite, $B_1(z)$	bip. bridgeless, $B_3(z)$	$z/(1 - z(1 + M))^2$	$z(1 + M)^2$
bipartite, $B_1(z)$	bip. nonsep., $B_4(z)$	$z(1 + M)^2$	–
bip. nonsep., $B_4(z)$	bip. ns. smpl, $B_5(z)$	$z(1 + M)$	–
singular tri., $T_1(z)$	triang., $z + zT_2(z)$	$z(1 + M)^3$	–
triangulations, $T_2(z)$	irreducible tri., $T_3(z)$	$z(1 + M)^2$	–

Extension to more involved decompositions

- For maps : maps into loopless blocks, 2-connected maps into 3-connected blocks;
- For decorated maps : tree-rooted quadrangulations into simple blocks, Schnyder woods / 3-orientations / 2-orientations into irreducible blocks.

Critical window?

Phase transition very sharp => what if $u = 9/5 \pm \varepsilon(n)$?

- Block size results still hold if $u_n = 9/5 - \varepsilon(n)$, $\varepsilon^3 n \rightarrow \infty$;
- For $u_n = 9/5 + \varepsilon(n)$, this is the case as well: when $\varepsilon^3 n \rightarrow \infty$

$$L_{n,1} \sim 2.7648 \varepsilon^{-2} \ln(\varepsilon^3 n)$$

(analogous to [Bollobás 1984]'s result for Erdős-Rényi graphs!);

- Results exist for scaling limits in ER graphs [Addario-Berry, Broutin, Goldschmidt 2010], open question in our case.

Is there a critical window? If so, what is its width?

Thank you!